



Analysis of Soil Quality in a Deltaic Hydrocarbon Polluted Environment Niger Delta Nigeria

Chukwu Okeah, G. O.^{1*}, Ushie, Lawrence¹ and Oyegun, Charles Uwadiae¹

¹*Department of Geography and Environmental Management, University of Port Harcourt, P.M.B. 5323, Port Harcourt, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author COGO was instrumental to the collection and analysis of the data for this study and also took active part in the drafting. Author UL initiated the research and was in charge of every logistics the work required he was also a part of the drafting of this work, he made the first manuscript of this work and finally author OCU gave a final touch on this work, He ensured that all the structures were done appropriately and proof read as well as edited and also made useful contribution to the overall study.

Article Information

Editor(s):

(1) Dr. Elmarie van der Watt, Department of Soil, Crop and Climate Sciences, University of the Free State, South Africa.

Reviewers:

(1) Ariech Mounira, Mohamed Boudiaf University, Algeria.

(2) Ronald Bartzatt, University of Nebraska, USA.

(3) Franco Cervellati, University of Ferrara, Italy.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/50709>

Original Research Article

Received 10 June 2019
Accepted 16 August 2019
Published 31 December 2019

ABSTRACT

Aim: The study was aimed to ascertain the damage done by an oil spill on the soils of Ogbia area in Bayelsa State, Niger Delta, Nigeria.

Study Design: The study adopted the experimental research design which entailed the use of field measurements and a control site. Three communities which are Elebele, Imiringi and Otuasaga were used for the study. At each of the sample locations, three random spots were augered at two depth-levels (Top Sample (T), 0 – 15 cm; Bottom Sample (B), 15- 30 cm), with the aid of an auger to collect the samples for laboratory analysis. The parameters of interest to the study are TPH, THC, Organic Matter, THF, THB, pH, sand, silt clay and soil texture and these were analysed using standard techniques as recommended by DPR.

Results: The study revealed that there is a noticeable effect of oil exploitation activities on soil quality within the study area, In the case of THC, the result revealed that the level of THC in the sampled communities was higher than that of the non spill site, hence the presence of hydrocarbon

*Corresponding author: Email: gift.chukwuokeah@uniport.edu.ng, giftchukwuokeah@yahoo.com;

which has caused the pollution of the soil. Organic matter content of the soil also reveals that the non-polluted site has more organic content than the sampled communities.

Conclusion: The study revealed that there is a statistically significant difference in soil quality of the selected communities and that of the non-spilt site. The study, therefore, recommended complete and total remediation of the soils in the area, as this will enhance the soil for increased food production.

Keywords: Soil; hydrocarbon; polluted; deltaic; environment.

1. INTRODUCTION

Over the years there have been a rising demand for oil and the dependence on oil revenue has greatly increased. This increase has brought an increase in the economic development of Nigeria, but the situation has led to an increase in environmental, degradation, pollution and contamination of the soils in the Niger Delta. This has arisen from the breakdown of oil pipelines and wells as well as petroleum distribution processes. Oil spills on land have brought about a reduction in soil fertility which has affected food production [1,2].

While the benefit of the oil industry is not in doubt, it's explorative, developmental and production processes generate a lot of waste amongst which are of "drilling cuttings, drilling fluids, produced water, sludge, completion and work over fluids, trace metals, heat waste and oxides of carbon, Sulphur and nitrogen" [3]. Fluids and cuttings arising from oil drilling are the largest waste sources during exploration. It is speculated that at least two barrels of fluids and cuttings are generated per foot of a typical well drilled [3].

Several studies have revealed that oil pollution affects the quality of soils; in the study conducted by Bada and Olarinre [4], revealed that leaves had more heavy metal content in plants than stems and roots.

Furthermore, Oyem and Oyem [5], in Ugborodo community on oil spillage impacts on the soil physico-chemical properties. Arising from oil spillage, it was observed that among the four sampled communities parameters measured indicated high amount of hydrocarbon, hence a turn out low fertility, which will bring about low food production and its attendant effects on the sources of livelihood available to the people.

Ugboma, [6] in a separate study on showed that soil physical and chemical characteristics were affected by oil spill which resulted in soil fertility

decrease and crop productivity decline as well. This, therefore, provides evidence that oil spill affects soil quality which in turn affects the quantity of food production in an area.

The record of oil exploitation activities can be likened to the record of oil poison. This is because as oil exploration and exploitation commenced the next that happened "almost immediately was the three major causes of oil pollution namely; the impact of the seismic survey, gas flaring and oil spills" [7]. The consequences of oil exploitation on the environment with an emphasis on the soil quality informed the need for this study.

Through a large number of literature research, the harm of oil-polluted soil mainly includes the following aspects: Firstly, because of the small density, higher viscosity and lower emulsifying ability of petroleum, it is easy to be absorbed in soil surface, affecting the permeability and porosity of soil [8,9]; petroleum is rich in carbon and a small number of nitrogen compounds, so it can change the composition and structure of soil organic matter and impact the C/N, C/P, salinity, pH, EH and conductivity of soil [10].

The heavy metals (nickel and vanadium) in oil mixtures [11] and high concentrations of salt in oilfield output water can also damage the soil environment [12]. Secondly, microorganisms in the natural environment are quite abundant in healthy and clean soil. In normal situation, the microorganisms which can resist the oil pollution stress are not developed, while in contaminated soil, in order to adapt to this kind of environment, they can produce certain enzyme system and gradually form a dominant population with symbiotic or synergy effect [13].

A number of studies have shown that the hydrocarbon pollution can change the microbial population, the composition of the community structure and the enzyme system in soil, given priority to the inhibitory action [14,15]. Thirdly, it can impede the normal growth of crops such as

reduce the germination rate and fertility and decline the resistance to pests and diseases [16,17,18].

Moreover, the polycyclic aromatic hydrocarbons in petroleum chemicals have carcinogenic, mutagenic, teratogenic and other toxic effects. It can enter into the bodies of people and animals through breathing, skin contact and diet, degrading the normal function of livers and kidney etc, therefore causing a great threat to human's health. At last, the oil pollutants in the soil not only impact the pedosphere, but also the atmosphere and water sphere.

2. MATERIALS AND METHODS

2.1 The Study Area

The area of study is Imiringi, Otuasega and Elebele communities which are located within the Kolo Creek, which is located $4^{\circ}47'0''\text{N}$ – $6^{\circ}25'0''\text{E}$ within the lower Niger Delta. (See Fig. 1).

Ogbia is one amongst other local government areas in Bayelsa State, with the Head Quarters situated in the town called Ogbia which is located on the South of the area within Latitude $4^{\circ}45'00''\text{N}$ $6^{\circ}39'00''\text{E}$. It covers a total area of 695 km^2 . Kolo Creek oil and gas field as called by SPDC is located in Imiringi town. The name Kolo creek as called by SPDC, and is “named after Kolo Creek in Ogbia Local Government Area of Bayelsa State. The area is located about 10 km North-West of Yenagoa, which is the capital city of Bayelsa State. It is characterized by tropical rain forest and fresh water swamps that are usually flooded in the raining season”.

The study was conducted in the Kolo Creek area as called by Shell Development Company of Nigeria (SPDC) which is basically made up of three communities in Ogbia; there are Imiringi, Otuasega having and Elebele communities which were selected for the study because of the presence of oil wells in the communities. The study adopted the use of the experimental research design which enabled the researchers achieve the purpose of the study.




Fig. 1. Ogbia Area showing sampled communities

Since the interest of the study is on soil quality, soil samples were collected using a systematic sampling pattern (Tel and Hagarty, 1984). At each of the sample locations, three random spots were augered at two depth-levels (Top Sample (T), 0 – 15 cm; Bottom Sample (B), 15- 30 cm), with the aid of an auger [19]. Also, at each of the sample locations and soil depth levels (T or B), the soil samples were bulked together to give a composite sample. The soil samples from different sample locations and soil depth levels were on each occasion collected in polythene bags and labeled accordingly and sent to the laboratory for analysis. Two hypotheses were also put forth to guide this study.

2.2 Analytical Procedure for Soil Quality Analysis

2.2.1 Total heterotrophic bacteria

A sample of water with a mass of 1 gram is weighed into 9ml sterile diluents (0.85% NaCl) under aseptic condition. It is shaken vigorously to homogenize and serially diluted. Then 0.1ml aliquot of the inoculums is collected using a sterile pipette, inoculated on Nutrient Agar (NA) surface. The inoculums are spread evenly with sterile hockey stick (Bent rod). Plates are incubated at 28°C for 24 hours. Thereafter, colonies are counted to obtain colony forming unit (cfu) value per ml of water sample. Distinct colonies are picked and streaked or subculture on freshly prepared Nutrient Agar medium to obtain pure culture after 24 hours incubation at 37°C. The pure culture is gram stained for microscopic examination. It is also used to carry out biochemical tests for characterization and identification of the isolates.

2.2.2 Total petroleum hydrocarbon

This analysis shall be done using gas chromatography/ flame ionization detector method. Methods 3580A and 3550C describe the procedure used for extracting non-volatile and semi-volatile organic compounds from solids such as soils, sludge and wastes, while method 8000 describes the determinative steps employed in the GC/FID of sample extracts obtained using the methods described above.

2.2.3 Soil pH determination

The pH of soil was done with the aid of a pH meter.

3. RESULTS

The results of the laboratory analysis of the soil samples collected from the selected communities are presented in the Table 1.

4. DISCUSSION

As seen in Table 1, Total Petroleum Hydrocarbon in soils of the selected communities ranges between 132.51 and 1,732.50, with Otuasega community accounting for the lowest which is 132.51 mg/kg and Elebele community accounting for 1,732.50 mg/kg which is the highest, although the result shows that total petroleum hydrocarbon value differs significantly from one location to another.

Total Hydrocarbon Content in soils of the selected communities ranges between 223.7 mg/kg and 2246.0 mg/kg, with Otuasega community accounting for the lowest which is 223.7 mg/kg and Elebele community accounting for 2246 mg/kg which is the highest; although the result shows that total hydrocarbon content value in soils differs significantly from one location to another.

Organic matter in soils of the selected communities ranges between 7.440 and 14.890, with Elebele community accounting for the highest, with an organic content value of 14.890. This was followed by Otuasega community with an organic content value of 11.280 and Imiringi community with an organic content value of 7.440.

Total Heterotrophic Fungi in soils of the selected communities ranges between 0.2 and 1.2, with Imiringi community accounting for the highest, with a THF value of 1.2. The two other sampled community's Otuasega and Elebele had the same THF value of 0.2 each.

Total Heterotrophic Bacteria in soils of the selected communities ranges between 1.0 and 3.2, with Imiringi community accounting for the highest, with a THB value of 3.2. This was followed by Otuasega community with a THB value of 1.2 and Elebele community with a THB value of 1.2.

pH in soils of the selected communities ranges between 5.30 and 6.50, with Elebele community accounting for the highest, with a pH value of 6.50. This was followed by Imiringi community with a pH value of 6.30 and Imiringi community with a pH value of 5.30.

Table 1. Result of physico-chemical/microbiological analysis of soil

Parameters	Imiringi	Elebele	Otuasega
TPH(mg/kg)	178.77	1,732.50	132.51
THC(mg/kg)	277.2	2246.0	223.7
Organic matter (%)	7.440	14.890	11.280
THF(cfu/g)x10 ³	1.2	0.2	0.2
THB(cfu/g)x10 ³	3.2	1.0	1.2
P ^H	6.30	6.50	5.30
Sand (%)	11.2	12.7	11.9
Silt (%)	31.6	33.4	32.8
Clay (%)	57.2	53.9	55.3
PSD/Texture	Silty clay	Silty clay	Silty clay

Source: Researchers field work (2015)

Table 2. Comparism of soil quality parameters of polluted sites with non polluted soil

Parameters	Imiringi	Elebele	Otuasega	Non polluted site
TPH(mg/kg)	178.77	1,732.50	132.51	20.18
THC(mg/kg)	277.2	2246.0	223.7	36.0
Organic matter (%)	7.440	14.890	11.280	18.20
THF(cfu/g)x10 ³	1.2	0.2	0.2	0.5
THB(cfu/g)x10 ³	3.2	1.0	1.2	2.2
P ^H	6.30	6.50	5.30	7.10
Sand (%)	11.2	12.7	11.9	13.8
Silt (%)	31.6	33.4	32.8	35.2
Clay (%)	57.2	53.9	55.3	51
PSD/Texture	Silty clay	Silty clay	Silty clay	Silty clay

Source: Researchers field work (2018)

Table 3. Chi-square analysis for the difference in soil quality between Imiringi (Polluted site) and non-polluted site

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
178.77	20.18	158.9	25150.79	1246.3
277.2	36.0	241.2	58177.4	1616.0
7.440	18.20	-10.76	115.78	6.36
1.2	0.5	0.7	0.49	0.98
3.2	2.2	1	1	0.45
6.30	7.10	-0.8	0.64	0.090
11.2	13.8	-2.6	45.69	3.31
31.6	35.2	-3.6	12.96	0.368
57.2	51	8.2	67.24	1.32
				$X^2 = 2875.12$

$$df = (9-1) (2-1)$$

$$(9-1) (2-1)$$

$$8 \times 1 = 8$$

Therefore we conclude that X^2 calculated value is 2875.12 while the critical value at 10 degree of freedom and 95% significant level is 15.51

The calculated X^2 calculated value 2875.12 is greater than the critical value of 15.51 we, therefore, reject the null hypothesis which states that there is no statistically significant difference in soil quality of Imiringi and that of the non polluted site and accepts the alternate hypothesis which states that there is a statistically significant difference in soil quality in Imiringi and that of the non polluted site

Sand level in soils of the selected communities ranges between 11.2% and 12.7% with Elebele community accounting for the highest, with sand

% value of 12.7%. This was followed by Otuasega community with sand % value of 11.9% and Imiringi community with sand % value

of 11.2%.Silt as observed to range between 31.6 and 33.4. The highest was recorded in Elebele with a percentage value of 33.4% silt; Otuasega had 32.8% silt and Imiringi 31.6% silt.

Clay ranged between 53.9 and 57.2 with the highest percentage value of 57.2% recorded in Imiringi. This was followed by Otuasega with clay percentage value of 55.3% and Elebele with a

clay percentage value of 53.9%.The texture of the soil as shown revealed that the three community soils are silty clay soil.

Table 2 revealed that total petroleum hydrocarbon level in the three sampled communities is above that of the non polluted site with Elebele community having more concentration of total petroleum hydrocarbon

Table 4. Chi-square analysis for difference in soil quality between Elebele (Polluted site) and non-polluted site

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
1,732.50	20.18	1712.32	2932.01	145.3
2246.0	36.0	2210	4884100	135669.4
14.890	18.20	-3.31	10.96	0.602
0.2	0.5	-0.3	0.09	0.18
1.0	2.2	-1.2	1.44	0.65
6.50	7.10	-0.6	0.36	0.05
12.7	13.8	-1.1	1.21	0.088
33.4	35.2	-1.8	3.24	0.092
53.9	51	2.9	8.41	0.165
				$X^2 = 135,816.5$

$$df = (9-1) (2-1) \\ (9-1) (2-1) \\ 8 \times 1 = 8$$

Therefore we conclude that X^2 calculated value is 135,816.5while the critical value at 10 degree of freedom and 95% significant level is 15.51

The calculated X^2 calculated value 135,816.5is greater than the critical value of 15.51 we therefore reject the null hypothesis which states that there is no statistically significant difference in soil quality of Elebele and that of the non polluted site and accept the alternate hypothesis which states that there is a statistically significant difference in soil quality in Elebele and that of the non polluted site

Table 5. Chi-Square analysis for the difference in soil quality between Otuasega (Polluted site) and non-polluted site

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
132.51	20.18	112.33	12618.03	625.3
223.7	36.0	187.7	35231.3	978.65
11.280	18.20	-6.92	47.89	2.63
0.2	0.5	-0.3	0.09	0.18
1.2	2.2	-1	1	0.45
5.30	7.10	-1.8	3.24	0.456
11.9	13.8	-1.9	3.61	0.263
32.8	35.2	-2.4	5.76	0.164
55.3	51	4.3	18.49	0.363
				$X^2 = 1608.456$

$$df = (9-1) (2-1) \\ (9-1) (2-1) \\ 8 \times 1 = 8$$

Therefore we conclude that X^2 calculated value is 1608.456while the critical value at 10 degree of freedom and 95% significant level is 15.51

The calculated X^2 calculated value 1608.456is greater than the critical value of 15.51 we therefore reject the null hypothesis which states that there is no statistically significant difference in soil quality of Otuasega and that of the non polluted site and accept the alternate hypothesis which states that there is a statistically significant difference in soil quality in Otuasega and that of the non polluted site

Table 6. One way (ANOVA) for variation in soil quality in sampled communities in Ogbia

Source of variation	ANOVA					
	SS	Df	MS	F	P-value	F crit
Between Groups	948295.4	2	474147.7	1.804708	0.186144	3.402826
Within Groups	6305476	24	262728.2			
Total	7253771	26				

than the other communities. The result here shows that there is a noticeable effect of oil exploitation activities of soil quality within the study area.

In the case of THC, the result revealed that the level of THC in the sampled communities was higher than that of the non-spill site, hence the presence of hydrocarbon which has caused the pollution of the soil.

The organic matter content of the soil also reveals that the non-polluted site has more organic content than the selected communities. This implies that the presence of oil in the soil due to pollution has affected the organic content of the soil hence bringing a reduction in its amount when compared with the non-polluted site.

4.1 Hypothesis Testing

In this section effort was made towards testing the stated study hypotheses with the aid of the Chi-Square statistical tool. Since two hypotheses were postulated for this study both were also tested in this section.

4.2 Hypothesis Two

The second hypothesis of the study, states that there is no statistically significant variation in the extent of soil damage amongst sampled communities in Ogbia.

The Table 6 shows the variation in the extent of soil damage amongst sampled communities in Ogbia.

From the Table 6, it is revealed that calculated F value for the analysis is 1.804708 while the critical value is 3.402826. Since the calculated F value of 1.804708 is less than the critical value of 3.402826 at F_{26}^2 degree of freedom, the implication of this is that the null hypothesis H_0 of no significant difference is accepted while rejecting the alternate H_1 which states that there is a statistically significant variation in the extent of soil damage amongst selected communities.

Arising from the above it is pertinent to state that the study has revealed that there is no statistically significant variation in soil quality of the selected communities.

5. CONCLUSION

The study concludes that the level of damage done by oil spill on the soils and its spatial variation in Ogbia, the study therefore revealed that total petroleum hydrocarbon level in the three sampled communities is above that of the non polluted site with Elebele community having more concentration of total petroleum hydrocarbon than the other communities. The result here shows that there is a noticeable effect of oil exploitation activities on soil quality within the study area. For THC, the result revealed that the level of THC in the sampled communities was higher than that of the non spill site, hence the presence of hydrocarbon which have caused the pollution of the soil. The organic matter content of the soil also reveals that the non polluted site has more organic content than the sampled communities. This implies that the presence of oil in the soil due to pollution has affected the organic content of the soil hence bringing a reduction in its amount when compared with the no polluted site.

Statistically, the study revealed that there is a statistically significant difference in soil quality of the sampled communities and that of the non spilled site, but on specific parameter, it was found that total petroleum hydrocarbon and total hydrocarbon content were higher in the selected communities than the non oil spilled site. The study further revealed that;

1. There is a noticeable effect of oil exploitation activities on soil quality within the study area.
2. There is also a noticeable decline in the crop production arising from soil pollution occasioned by oil exploitation.
3. There is a clear cut evidence of the state of the soil in the polluted sites as compared to the non polluted which is an

indication of the state of the soil having been ravaged by the detrimental effect of oil exploratory activities in the area.

4. Total petroleum hydrocarbon level in the three sampled communities is above that of the non polluted site with Elebele community having more concentration of total petroleum hydrocarbon than the other communities.
5. The study revealed that there is a statistically significant difference in soil quality of the selected communities and that of the non spilled site.

On the basis of the above, the study recommended a complete and total remediation of the soils in the area, as this will enhance the soil for increased food production.

CONSENT

Before the commencement of the research the communities were duly informed of the research which they obliged and provided a guard to assist the researchers during the period.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sparrow SD, Sparrow EB. Microbial biomass and activity in a subarctic soil ten years after crude oil spills. *Journal of Environmental Quality*. 1988;17:304–309.
2. Racine Ch. H. Long-term recovery of vegetation on two experimental crude oil spills in interior Alaska black spruce taiga. *Canadian Journal of Botany*. 1993;72: 1171–1177.
3. Nwilo PC, Badejo OT. Oil spills problem and management in the Niger Delta International Oil Spills, Conference Monitoring. 2005;2.
4. Bada BS, Olarinre TA. Characteristics of soils and heavy metal content of vegetation in oil spill impacted land in Nigeria. *Proceedings of the Annual International Conference on Soils, Sediments, Water and Energy*. 2012;17(2). Available: http://scholarworks.umass.edu/s_oils_proceedings
5. Oyem, Isama Lawrence Rank, Oyem, Isama Lawrence. Effects of crude oil spillage on soil physico-chemical properties in Ugborodo Community. *International Journal of Modern Engineering Research (IJMER)*. 2013;3(6): 3336-3342. Available: www.ijmer.com [ISSN: 2249-6645] Available: www.ijmer.com
6. Ugboma PP. Effects of oil spillage on soil fertility in Udu Local Government Area in Delta State. *International Journal of science and Technology, Ethiopia*. 2014;3 (3):47-56. [ISSN: 2225-8590] [ISSN 2227- 5452] DOI: <http://dx.doi.org/10.4314/stech.v3i3.5>
7. Pyagbara LS. The adverse impacts of oil pollution on the environment and wellbeing of a local indigenous community: The experience of the Ogoni. New York: Department of Economic and social Affairs, United Nations; 2007.
8. Wang B. Mechanism of growth and physiological response of three arbor species to oil contamination. Shaanxi: Northwest A&F University; 2014.
9. He LJ, Wei DZ, Zhang WQ. Research of microbial treatment of petroleum contaminated soil. *Advances in Environmental Science*. 1999;7(3):110-111 (in Chinese).
10. Li L, Liu M, Zhao JL. Current status and prospects of microbial remediation of soil pollution in oil fields in northern Shaanxi. *Chinese Journal of Soil Science*. 2011;42 (4):1011-1013.
11. Saadat S, Mirkhani R, Mohebi A. Study on phytoremediation of soils polluted with heavy metals and oil pollutants in agricultural lands affected by Persian Gulf War (Khouzestan, fars, kohgiluyeh & boyrahmad and boushehr provinces); 2014.
12. Efsun DF, Olcay Hüseyin S. Variations of soil enzyme activities in petroleum-hydrocarbon contaminated soil *International Biodeterioration & Biodegradation*. 2015;268-275.
13. Chiara A, Rosario M, Flavia T. Bioremediation of diesel oil in a co-contaminated soil by bioaugmentation with a microbial formula tailored with native strains elected for heavy metals resistance. *Science of the Total Environment*. 2009;407(8):3024-3032.
14. Deng RY. Microbiological monitoring and evaluation of compound pollution of petroleum and heavy metal in slated soils.

- Shandong: Shan Dong University, China; 2014.
15. Uzoije AP, Agunwamba JC. Physiochemical properties of soil in relation to varying rates of crude oil pollution. *Journal of Environmental Science and Technology*. 2011;4(3):313-323.
 16. Xu YH, Lu M. Bioremediation of crude oil-contaminated soil: Comparison of different biostimulation and bioaugmentation treatments. *Hazardous Materials*. 2010;183:398-401.
 17. Zhu H, Liu K, Yang X. Sedimentary controls on the sequence stratigraphic architecture in intracratonic basins: An example from the Lower Permian Shanxi Formation, Ordos Basin, Northern China. *Marine and Petroleum Geology*. 2013;45:42-54.
 18. Shan BQ, Zhang YT, Cao QL. Growth responses of six leguminous plants adaptable in Northern Shaanxi to petroleum contaminated soil. *Environmental Science*. 2014;35:1125-1130.
 19. Department of Petroleum Resources (DPR). Environmental Guidelines and Standards for the Petroleum Industry in Nigeria (EGASPIN). 2002;59–66.

© 2019 Chukwu-Okeah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/50709>