



Exploring the Factors Associated with Birth Asphyxia among the Newborn Infants at a Rural Hospital in Bangladesh

Shabnam Mostari¹, Md. Razwan Hasan Khan Chowdhury²,
Mohammad Rocky Khan Chowdhury³, Dioneia Motta Monte-Serrat⁴,
Masoud Mohammadnezhad⁵ and Russell Kabir^{6*}

¹The Obstetrical and Gynecological Society of Bangladesh, Dhaka, Bangladesh.

²Department of Sociology, First Capital University, Chuadanga, Bangladesh.

³Department of Public Health, First Capital University, Chuadanga, Bangladesh.

⁴Department of Health and Education, Universidade de Ribeirao Preto, Sao Paulo State, Brazil.

⁵School of Public Health & Primary Care, Fiji National University, Fiji.

⁶School of Allied Health, Anglia Ruskin University, Essex, UK.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRNH/2018/45726

Editor(s):

(1) Dr. Abdel-Hady El-Gilany, Professor, Department of Community Medicine, Mansoura University, Egypt.

(2) Dr. Carla Maria Ferreira Guerreiro da Silva Mendes, Nurse Director at Centro Hospitalar de Setúbal, Portugal.

(3) Dr. Fernanda Jorge Guimaraes, Professor, Universidade Federal de Pernambuco, Centro Academico de Vitoria, Rua Alto do Reservatorio, Brasil.

Reviewers:

(1) Elkenah Ndie Chubike, National Open University of Nigeria, Nigeria.

(2) K. Ramesh Kumar, S.V.S.Medical College and Hospital, K.L.N.university of Health Sciences, India.

(3) Mario Ganau, University of Toronto, Canada.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/45726>

Original Research Article

Received 25 September 2018

Accepted 04 January 2019

Published 18 January 2019

ABSTRACT

Background: Birth asphyxia is a critical problem to address within the context of public health. There are many reasons a baby may not be able to take in enough oxygen before, during, or just after birth.

Objective: The objective of this research is to determine the risk factors that are associated with birth asphyxia in the Matlab, Hospital Bangladesh.

Methods: This is a case-control study. Cases were selected from the hospital records. A group of

*Corresponding author: Email: russell.kabir@anglia.ac.uk, russellkabir@live.co.uk;

94 mothers who gave live births resulting in asphyxiated new-borns during 24th October 2006 to 20th August 2008 at the labour ward of Matlab hospital, Bangladesh.

Results: Result indicates the largest group of mothers with Asphyxiated new-born between <25 years old, was 61 (35.5%) while in the control group it was 111 (64.5%). Foetal presentation was associated with birth asphyxia in new-born (OR= 15.21; 95% CI: 1.841-125.67; p= 0.001). The study shows that infant with birth weight <2500g had 3.49 times greater risk of developing birth asphyxia compared to the infant with birth weight >2500 g (CI: 1.79-6.78, p = 0.00).

Conclusions: A future community based study with larger sample size is necessary to find the temporal relationship between Birth asphyxia and cord around the neck, Gestational age at birth less than 37 weeks and prolong first stage of labour.

Keywords: Birth asphyxia; newborns; Bangladesh; rural; hospital.

1. BACKGROUND

Birth asphyxia is a critical problem to address within the context of public health. According to the World Health Organization (WHO) birth asphyxia is defined as "failure to initiate and sustain breathing at birth" and based on Apgar score as an Apgar score of <7 at one minute of life [1]. Each year, globally, it results in many neonatal deaths and neurological disability especially in low and middle-income countries after infections and preterm births. It is the third major cause of neonatal death. Globally, birth asphyxia is estimated to account for 23% of the 4 million neonatal deaths [2] and 26% of the 3.2 million stillbirths each year [3,4]. According to WHO about 3% from 120 million infants born every year in developing countries suffers from birth asphyxia and estimated 900,000 die each year [5].

One of the present challenges is the lack of a gold standard for accurately defining birth asphyxia. Because of the same reason the incidence of birth asphyxia is difficult to quantify. Definitions of birth asphyxia designed for use in hospital-based settings require an evaluation of neonatal umbilical cord pH, Apgar scores, neurological clinical status, and markers of multisystem organ function, [6] and are not feasible for community settings [7]. In our study we have used Apgar score to determine the condition of Birth asphyxia. The risk factors for birth asphyxia in hospital-based and home-based may be similar. Birth asphyxia is a major neonatal health problem in Bangladesh and it is more common in rural areas. Because in rural areas most of the births are attended by untrained birth and antenatal risk factors are not identified. This is because pregnant mothers are not empowered to seek healthcare services [8]. Therefore risk factors are not diagnosed in time. When these deliveries are conducted at home by

untrained personnel- results in perinatal asphyxia in most of the situations. Birth asphyxia is the main causes of neonatal mortality especially in low income countries and there is a dearth of publications on the incidence and risk factors of severe birth asphyxia in Bangladesh. On this backdrop, the purpose of this research is to explore the factors responsible for birth asphyxia in the Matlab Hospital, Bangladesh.

2. METHODS AND DATA

This is a retrospective case –control research design. The study populations were the mothers who gave live birth at Matlab hospital. In a series of 282 mothers were selected, 94 mothers were given live births resulting in asphyxiated new-born and other 188 mothers gave birth to healthy new-born. Apgar score was used to diagnose birth asphyxia in the present study. Apgar score of less than < 7 at 1 minute of birth was included. Apgar scores less than 7 were considered as cases and Apgar scores 7 or more at 1 minutes of birth were in the control group.

2.1 Selection of Cases

Cases were selected from the hospital records. A group of 94 mothers who gave live births resulting in asphyxiated new-borns during 24th October 2006 to 20th August 2008 at the labour ward of Matlab hospital, Bangladesh.

2.2 Selection of Control

Control is the group of 188 mothers who gave live birth resulting in non-asphyxiated new-borns at the same hospital at the same period of time of cases and whose Apgar score more than 7 at 1 minute of birth. To avoid misclassification bias, controls were selected from the groups of newborn who's Apgar score more than 7 at one minute of birth. Two cases of control group

compared to one case of study group by selecting the sample case which is prior and after to a sample case in the study group.

2.3 Sample Size Determination

The sample size was determined by using Epi-info version 6. The probability that if the two samples differ this reflects a true difference in the two populations (confidence level or $1-\alpha$) = 0.05 Probability that if the two populations differ, the two samples will show a 'significant' difference (power or $1-\beta$) = 75.

Proportion of controls with exposure = 0.5; Odd Ratio= 2; Control: Case= 2:1

By using Epi-info version, I have found my cases that was 94;

So the control was $94 \times 2 = 188$;

And total sample size was $188+92= 282$

Inclusion criteria- (i) The mother who gave live birth having Apgar score was less than < 7 at 1 minute of birth (ii) The mothers with 28 weeks gestation or more (iii) Mother who deliver at Matlab hospital.

Exclusion criteria- (i) Children with congenital anomalies and genetic disorders were excluded from the study.

Data were collected by reviewing the hospital records of all the cases and the selected controls by a nurse who works in the Matlab Hospital. Additionally, the delivery records of expectant mothers resulting in the live births have been also included in this study. The nurse was trained properly before data collection and was blinded about the objectives. The data was recorded to the specific data collection form, systematically designed for computer coding and analysis. Information obtained from the cases and controls were coded and entered twice into the microcomputer to avoid the mistake. Data were analysed using Statistical Package for Social Sciences (SPSS) ver. 16. Quality and reliability of collected data were re-checked. Following the WHO and Bangladesh Medical Research Council (BMRC) guidelines of ethical consideration, and the informed consent was taken before the data collection process.

3. RESULTS

Table 1 shows the general characteristics of the subjects of case and control groups including the OR, with the association of birth asphyxia. This

result indicates the largest group of the mother with Asphyxiated new-born between <25 years old, was 61 (35.5%) while in the control group it was 111 (64.5%). On the other hand, the second largest group age between 26-35 years old was in the case group of 21(33.3%) and the control group were 42 (66.7%). The third largest group below 20 years old or lower being in the case group were 11 (26.8%) and control group were 30 (73.2%). The smallest group found in oldest mother with ages 36 years old or higher at 1(20%) in case group and control group were 4(80%). However maternal age below 25 years (OR= 1.49; 95% CI: 0.70-3.19; $p= 0.29$) and age between 26-35 years (OR= 1.27; 95% CI: 0.54-3; $p= 0.57$) was not statistically significant association between maternal age and asphyxia in new-born. Having one to three children in the family had OR of 0.80 and more than three children had OR of 0.59 which were not statistically significant (95% CI: 0.48-1.32; $p= 0.39$) for one to three children and (95% CI: 0.11-3.03; $p= 0.52$) for more than three children. From the statistical significance test result, it suggests that maternal Height less than 145 cm had 2.22 times more risk of getting birth asphyxia on her neonates compared with mother more than 145cm height which was statistically significant (OR= 2.22 ; 95% CI: 1.14-4.31; $p= 0.01$).

Table 2 shows, 41.5% and 18.1% of newborn both cases and controls had a birth weight less than 2500 gm while 58.5% and 82% of newborn both cases and controls had birth weight more than 2500 gm. The statistical significant analysis result suggests that low birth weight was associated with birth asphyxia in new-born compared with newborn more than 2500gm (OR= 3.21; 95% CI: 1.84-5.58; $p= 0.00$). It is found that gestational age 37-41 weeks (OR= 0.45; 95% CI: 0.20-1; $p= 0.057$) and >42 weeks (OR= 0.897; 95.0% CI: 0.32-2.48; $p= 0.83$) are not statistically significant in compared with gestational age <37 weeks at births. The head presentation of foetal are common in both mothers with asphyxiated new-born and non-asphyxiated new-born which consist of 92.0% and 99.4% of the sample. Whereas the breech/malpresentation of foetal is not common and only consist of 8.0% and 0.6% of the sample respectively. The logistic test result shows that foetal presentation was associated with birth asphyxia in new-born (OR= 15.21; 95% CI: 1.841-125.67; $p= 0.001$).

Table 3 shows that about 28.4% of the mother with asphyxiated new-born had Bad obstetric

history during delivery while 71.6% mother with asphyxiated new-born had no Bad obstetric history. About 11.4% of the mother with asphyxiated new-born had the history of Premature Labour while 88.6% mother with asphyxiated new-born had no such history. In contrast to mothers with non-asphyxiated new-born, 3.9% had the history of Premature Labour while 96.1% had no such history. The test result suggests that a history of Premature Labour was significantly associated with Birth Asphyxia in newborn (OR = 0.319; 95% CI: 1.149-8.533; p = 0.020). Also the results that the time taken in the Prolong second stage of delivery was associated with birth asphyxia in new-born (OR = 4.943; 95% CI= 1.661-14.71; p= 0.002) and, it suggests that the history of Meconium Stained Liquor were significantly associated with Birth Asphyxia in newborn (OR = 5.372; 95% CI: 2.479-11.64; p = 0.00).

minimum of four visits in antenatal care program and 34% of mothers who complete the three visits of antenatal care program and 19.1% of the mother complete 0-2 visit of antenatal care program. Similar result found in mother with non-asphyxiated new-born, about 54.3% complete the program, 26.6% mothers complete the three visits of antenatal care program and 19.1% of mother who complete 0-2 visit of antenatal care program. No statistically significant association was found between 0-2 visits of antenatal care and Birth Asphyxia compared to mother who visits total 4 antenatal visits of antenatal care program (OR= 1.15, 95% CI: .60-2.25, p= 0.66). However, Birth Asphyxia was not significantly associated with mothers who have completed the 3 visits of antenatal care program (OR= 1.48, 95% CI: .84-2.61, p=.17). The result also indicates that cytomis induction is not associated with birth asphyxia in new-born (OR = 1.11, 95% CI: 0.32-3.82, p = 0.863).

Table 4 shows there are about 46.8% of mothers with asphyxiated new-born who complete the

Table 1. Distribution of Socio demographic and maternal factor by case and control including OR and p-value

Categories	Cases (n= 94) No (%)	Controls (n= 188) No (%)	OR (95% CI)	p-value
Maternal age (Years)				
<25	45(48.0%)	92(49.0%)	1	
26-35	42(45.0%)	79(42.0%)	1.1(0.65-1.8)	0.75
>35	7(7.4%)	17(9.0%)	0.84(0.32-2.18)	0.72
Parity				
No parity	47(50.0%)	83(44.1%)	1	
1-3	45(47.9%)	99(52.7%)	0.80(0.48-1.32)	0.39
>3	2(2.1%)	6(3.2%)	0.59(0.11-3.03)	0.52
BMI				
<25	73(83%)	155(87.0%)	1	
>25	15(17%)	23(13%)	1.38(0.68-2.8)	0.36
Mother Height				
<145cm	21(24%)	22(12.4%)	2.22(1.14-4.31)	0.018*
>145cm	67(76.1%)	156(87.6%)	1	
Mother weight				
<50kg	55(59.8%)	86(47.3%)	1.65(0.99-2.75)	0.051*
>50kg	37(40.2%)	96(52.7%)	1	
Education				
No education	21(22.3%)	29(15.4%)	1	
Primary	25(26.6%)	58(31%)	0.59(0.28-1.23)	0.21
Secondary	48(51%)	101(53.7%)	0.65(0.34-1.26)	0.16
Asset index				
1(Poorest)	13(14.1%)	21(11.2%)	1	
2	20(21.7%)	35(18.6%)	0.92(0.38-2.23)	0.85
3	18(19.6%)	38(20.2%)	0.76(0.31-1.86)	0.55
4	16(17.4%)	40(21.3%)	0.64(0.26-1.59)	0.34
5(Richest)	25(27.2%)	54(28.7%)	0.74(0.32-1.73)	0.49

Table 2. Distribution of Foetal factors by case and control including OR and p-value

Categories	Cases (n= 94) No (%)	Controls (n= 188) No (%)	OR (95% CI)	p-value
Birth weight of infants (g)				
<2500gm	39(41.5%)	34(18.1%)	3.21(1.84-5.58)	0.00*
>2500gm	55(58.5%)	154(81.9%)	1	
Gestational age at birth (weeks)				
<37 weeks	13(13.8%)	14(7.4%)	1	
37-42 weeks	66(70.0%)	156(83%)	0.45(0.20-1)	0.057*
>42weeks	15(16.0%)	18(9.6%)	0.89(0.32-2.48)	0.83
Sex of infants				
Male	55(58.5%)	110(58.5%)	1	
Female	39(41.5%)	78(41.5%)	1(0.6-1.6)	1
Mal-presentation				
Present	7(8.0%)	1(0.6%)	15.21(1.84-125.67)	0.001*
Absent	81(92%)	176(99.4%)	1	
Cord around the neck				
Present	20(22.7%)	24(13.5%)	1.89(0.97-3.64)	0.056*
Absent	68(77.3%)	154(86.5%)	1	

Table 3. Distribution of deliveries related factor by case and control including OR and p-value

Categories	Cases (n= 94) No (%)	Controls (n=188) No (%)	OR (95% CI)	p value
Bad obstetric history:				
- Present	25 (28.4)	25 (28.4)	0.96	0.89
- Absent	63 (71.6)	63 (71.6)	(0.547-1.69)	1
Leaking Membrane:				
- Present	4 (4.5)	18 (10.1)	0.42	0.12
- Absent	84 (95.5)	160 (89.9)	(0.13-1.29)	1
Premature rupture of membrane:				
- Present	9 (10.2)	13(7.3)	1.44	0.41
- Absent	79 (89.8)	165(92.7)	(0.59-3.52)	1
Premature labor				
- Present	10 (11.4)	7 (3.9)	0.31	0.02*
- Absent	78(88.6)	171(96.1)	(1.14-8.53)	1
Prolong 1st stage				
-Present	15 (17.0)	25(14.0)	1.25	0.51
-Absent	73 (83.0)	153(86 .0)	(0.62-2.59)	1
Prolong 2nd Stage				
-Present	11 (12.5)	5(2.8)	4.94	0.00*
-Absent	77 (87.5)	173(97.2)	(1.66-14.71)	1
APH*				
-Present	1 (1.1)	1 (.6)	2.03	0.55
-Absent	87 (98.9)	177 (99.4)	(0.12-32.9)	
Meconium Stained Liquor				
-Present	23(26.1%)	65(73.9%)	5.37	0.00*
-Absent	11(6.2%)	167(93.8%)	(2.47-11.64)	1
Prolong labour				
-Present	15(17%)	7(3.9%)	5.02	0.00*
-Absent	73(83%)	171(96.1%)	(1.96-12.82)	1

Table 4. Distribution of maternal care received factor by case and control including OR and p-value

Categories	Case(n= 94) No (%)	Control (n= 188) No (%)	OR (95% CI)	p value
Antenatal care visits				
0-2 nd visits	18(19.1%)	36(19.1%)	1.15(0.59-2.25)	0.66
3 rd visits	32(34%)	50(26.6%)	1.48(0.84-2.61)	0.17
4 th visits	44(46.8%)	102(54.3%)	1	
Cytromis induction				
Yes	4(5.40%)	8(4.9%)	1	
No	70(94.6%)	156(95.1%)	1.11(0.32-3.82)	0.86

Table 5. Risk factor for Birth asphyxia in new-born

Factors	OR adjusted 95% CI	p Value	OR unadjusted 95% CI	p Value
Birth weight of newborn <2500gm	3.49 (1.79-6.78)	0.00	3.21 (1.84-5.58)	0.00*
Malpresentation	14.52 (1.59-132.62)	0.01	15.21 (1.84-125.67)	0.00*
Meconium stained liquor	6.80 (2.82-16.40)	0.00	5.37 (2.47-11.64)	0.00*
Prolong 2 nd Stage	4.96 (1.45-16.92)	0.01	4.94 (1.66-14.71)	0.00*
Premature labour	4.55 (1.62-12.74)	0.00	0.31 (1.15-8.53)	0.02*
Height of the mother	2.10 (1.13-3.89)	0.01	2.22 (1.14-4.31)	0.01*

Table 5 shows that the case and control status was fitted as the dependent variable into an Unconditional multiple logistic regression model with newborn weight (<2500gm,>2500gm), malpresentation, Meconium stained liquor, prolong the second stage of delivery and height of the mother (<145cm,>145 cm). The OR corrected for confounding factors were significantly associated with newborn weight, malpresentation, meconium stained liquor, prolong second stage of delivery and height of the mother. The infant with birth weight <2500g had 3.49 times greater risk of developing birth asphyxia compared to the infant with birth weight >2500g (CI: 1.79-6.78, p = 0.00). The malpresentation had 14.5 times greater risk of developing Birth asphyxia compared to the infant who's position was normal during delivery (CI: 1.59-132.62 p value = 0.018). Mother with the history of Meconium stained liquor had 6.8 times greater risk of developing birth asphyxia of the infants compared to the mother who had not such type of history (CI: 2.82-16.40, p = 0.00).The result in time study in prolong second stage of delivery shows that prolong 2nd stage had 4.9 times greater risk of developing birth

asphyxia of the infants compared to the mother who had not such type of history (CI: 1.45-16.92, p = 0.010).The study showed that mothers height <145cm had 2.1 times greater risk of developing birth asphyxia compared to the mother's height >145cm (CI: 1.13-3.89, p = 0.018).

4. DISCUSSION

In this Hospital based case-control study revealed that about 60% of male new-born had Birth asphyxia in both the cases and controls. Only 3% of both cases and controls had more than three children and rest had no children or up to three children that reflects well accepted family planning in the society. Several studies showed an association between parity and birth asphyxia [1-3]. This research revealed that there is no association between birth asphyxia and parity and similar to another study that obtained in India [4]. This study could not find any significant relationship between birth asphyxia and gestational age. And this study is similar to another study conducted at Jordan [5]. Again, this study could not find any association with maternal age, mother weight and mother BMI but

maternal weight was about to be associated. However, wide OR (95% CI: 0.99-2.75) indicated that with a larger sample size significant association could have been ascertained. A retrospective study conducted at Nepal have seen that maternal age between 18-35years was associated with Birth asphyxia [9]. This study could not find any significant relationship between birth asphyxia and gestational age and this study is similar to another study conducted at Jordan [5]. Moreover, there is no significant association found with mother age, mother weight and maternal BMI but the maternal weight was found to be associated. However, wide OR (95% CI: 0.99-2.75) indicated that with larger sample size significant association could have been ascertained. A retrospective study conducted at Nepal have seen that maternal age between 18-35years was associated with Birth asphyxia [9].

An infant with birth weight <2500g had 3.48 times greater risk of developing birth asphyxia compared to the infant with birth weight >2500gm. This finding is consistent with the findings of another study conducted at Nigeria and done over a three years period, showed that infants with low birth weight play a significant role in the occurrence of asphyxia [10]. A Retrospective case-control study conducted at Phramongkutklao hospital, Thailand have seen that fetal birth weight less than 2,500 grams had 2.5 times greater risk of developing birth asphyxia compared to the fetal birth weight more than 2500 grams [11]. But different from another hospital based study which was carried out in Jordan [5]. The study showed that mothers height <145cm had 2.1 times greater risk of developing birth asphyxia compared to the mothers' height >145cm. This is similar to a study conducted in Nepal [12].

Antenatal care from a trained provider is important to monitor the status of a pregnancy and to reduce the risk that is associated with the pregnancy and at delivery for the mother and child. The study findings show that majority of the mother had received four visits of antenatal care and only 19% of both cases and controls had received 0-2 visits of antenatal care and it showed that there was no significant relationship between birth asphyxia and antenatal care and the findings are a contrast to the findings in the Jordan study [5]. Around 29% of both cases and controls had bad obstetric history and this study also shows that, it is not associated with

birth asphyxia in new-born but this result is different from multiple hospital based studies [4,13,14]. In this study malpresentation is associated with birth asphyxia in new-born and this study is similar to another hospital based study which was conducted at Mulago hospital, Kampala, Uganda has seen that malpresentation had 6.32 times greater risk of developing Birth asphyxia compare to the normal presentation [15]. There was no significant difference in the premature rupture of the membrane of both the case and control group and it is not statistically significant but this study is different from other authors [1,13-16]. Prolonged rupture of the membrane may be associated with intrauterine infection resulting in birth asphyxia [1].

Meconium should always be considered a marker for foetal distress therefore there was a significant effect on the Apgar score of neonates [17]. Meconium-staining of the amniotic fluid is present in 9-14 per cent of all deliveries at the time of delivery [18]. In this study, meconium was strongly associated with Birth asphyxia in new-born and this study is similar to the other study [19-21]. A population based retrospective case-control study conducted at Sweden have seen that Meconium release had 4.1 times greater risk of developing birth asphyxia compared to the mother who did not have the history of Meconium release during delivery [22]. Coiling of the umbilical cord around the foetal neck is a common complication of labour, said to occur about once in every five deliveries [23]. In this study the association was borderline ($p=0.05$). Study conducted at London have found association between birth asphyxia and Cord around the neck [24]. Prolong second stage of labour is associated with birth asphyxia in new-born and this study is similar to another hospital based cohort study which was conducted at in New Delhi [10]. A retrospective study over a 3-year period from 1989 to 1991 was performed at a tertiary level, referral hospital have found association between birth asphyxia and prolonged second stage of labour [9].

4. LIMITATIONS

Potential limitation of the study could not be ignored. Firstly, the research relied on the Hospital records where history was taken from the attendant. The nurse who collected data was blinded of the objectives so; bias is unlikely in this regard. Secondly there was small sample size and as it is hospital-based study.

5. CONCLUSION AND RECOMMENDATION

Health education shows itself as a tool whose importance underlies some findings in this research. To Reduce Birth asphyxia requires the use of appropriate obstetric monitoring of pregnancy and labour for risk factors of Birth asphyxia and health education of the general population. Pregnancy related issues should be included in school curricula for adolescents. There is an immediate need for organization of regular workshops and seminars for TBAS, employees of public and private health institutions emphasising the need for early identification and prompt referral of complicated pregnancies and labour to appropriate health care institutions of efficient ambulance services and good network of roads. Further community-based study with larger sample size is necessary to find the temporal relationship between Birth asphyxia and cord around the neck, Gestational age at birth less than 37 weeks and prolong first stage of labour. More Hospital based prospective case-control study should be encouraged with larger sample size to identify the possible risk factors for Birth asphyxia. Implement the project to solve the problem from the factor found in this study and do the research to appraise the project of birth asphyxia in new-born problem solving project.

CONSENT AND ETHICAL APPROVAL

Following the WHO and Bangladesh Medical Research Council (BMRC) guidelines of ethical consideration, and the informed consent was taken before the data collection process.

FUNDING

There was no financial support received to conduct the research.

HIGHLIGHTS

1. Foetal presentation was associated with birth asphyxia in new-born babies.
2. Meconium was strongly associated with birth asphyxia
3. Infant with birth weight <2500 g had 3.48 times greater risk of developing birth asphyxia compared to the infant with birth weight >2500gm
4. There is no any significant relationship between birth asphyxia and gestational age

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Perinatal mortality. A listing of available information 4th ed. (WHO/FRH/MSM/95.7). Geneva, WHO Maternal Health and Safe Motherhood Programme; 1996.
2. Lawn JE, Cousens S, Zupan J. 4 million neonatal deaths: When? Where? Why? *Lancet*. 2005;365(9462):891-900. Mar 5-11
3. Lawn J, Shibuya K, Stein C. No cry at birth: Global estimates of intrapartum stillbirths and intrapartum-related neonatal deaths. *Bull Worl Healt Org*. 2005;83(6): 409-17.
4. Stanton C, Lawn JE, Rahman H, Wilczynska-Ketende K, Hill K. Stillbirth rates: Delivering estimates in 190 countries. *Lancet*. 2006;367(9521):1487-94.
5. World Health Organization. The World Health Report 1998—life in the 21st century: A vision for all. World Health Organization, Geneva, Switzerland; 1998.
6. SCORE A. Use and abuse of the Apgar score. *Pediatrics*. 1986;78(6).
7. Lawn JE, Manandhar A, Haws RA, Darmstadt GL. Reducing one million child deaths from birth asphyxia—a survey of health systems gaps and priorities. *Healt Res Policy Syst*. 2007;5:4.
8. Kabir R, Rahman S, Monte-Serrat DM, Arafat SY. Exploring the decision-making power of Bangladeshi women of reproductive age: Results from a national survey. *Sout East Asia Journ of Med Sci*. 2017;1(1):4-8.
9. Dongol SSJ, Shrestha S, Shakya A. Clinical profile of birth asphyxia in Dhulikhel Hospital: A retrospective study. *Jour of Nepal Paed Soc*. 2010;30(3):141-6.
10. World Health Organization. World Health Report 2003. Shaping the future. World Health Organization: Geneva; 2003.
11. Chayasak Pitsawong PP. Risk Factors associated with birth asphyxia in Phramongkutklao Hospital. *Thai Jour of Obs and Gyna*. 2011;19:165-71.
12. Nelson KB, Ellenberg JH. Apgar scores as predictors of chronic neurologic disability. *Pediatrics*. 1981;68(1):36-44.

13. World Health Organization. Neonatal and perinatal mortality: Country, regional and global estimates. WHO: Geneva; 2006.
14. Badawi N, Kurinczuk JJ, Keogh JM, et al. Intrapartum risk factors for newborn encephalopathy: The Western Australian case-control study. *BMJ*. 1998;317(7172): 1554-8.
15. Hall DR, Smith M, Smith J. Maternal factors contributing to asphyxia neonatorum. *Jour of Trop Ped*. 1996;42(4): 192-5.
16. Kinoti SN. Asphyxia of the newborn in east, central and southern Africa. *East Afr Med Jour*. 1993;70(7):422-33.
17. Bang AT, Bang RA. Diagnosis of causes of childhood deaths in developing countries by verbal autopsy: Suggested criteria. The search Team. *Bull Wor Heal Org*. 1992;70(4):499-507.
18. Lee AC, Mullany LC, Tielsch JM, et al. Risk factors for neonatal mortality due to birth asphyxia in southern Nepal: A prospective, community-based cohort study. *Pediatrics*. 2008;121(5):e1381-90.
19. Baqui AH, Darmstadt GL, Williams EK, et al. Rates, timing and causes of neonatal deaths in rural India: Implications for neonatal health programmes. *Bull Wor Heal Org*. 2006;84(9):706-13.
20. Chowdhury ME, Akhter HH, Chongsuvivatwong V, Geater AF. Neonatal mortality in rural Bangladesh: An exploratory study. *Jour of Healt Pop Nut*. 2005;23(1):16-24.
21. Bangladesh MOH & FW. National neonatal health strategy and guidelines for Bangladesh; 2009.
22. Milsom I, Ladfors L, Thiringer K, Niklasson A, Odeback A, Thornberg E. Influence of maternal, obstetric and fetal risk factors on the prevalence of birth asphyxia at term in a Swedish Urban population. *Acta Obs et Gyne Scand*. 2002;81(10):909-17.
23. National Institute of Population Research and Training MaA, and ORC Macro. Bangladesh Demographic and Health Survey; 2004.
24. Program for Appropriate Technology in Health (PATH) to save the children US. Jakarta I. Reducing birth asphyxia through the Bidan di Desa Program in Cirebon, Indonesia; 2006.

© 2018 Mostari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/45726>