

Current Journal of Applied Science and Technology

Volume 43, Issue 12, Page 199-204, 2024; Article no.CJAST.128455 ISSN: 2457-1024 (Past name: British Journal of Applied Science & Technology, Past ISSN: 2231-0843, NLM ID: 101664541)

# The Most Common Bacterial Infection among Iraqi Patients Suffering from Cholecystitis in AL-Anbar province: A Cross-sectional Observational Study

# Mohammed Ali Hussein <sup>a\*</sup>, Khalid Elhussein M. Idris <sup>b</sup> and Mutaman Ali A. Kehail <sup>b</sup>

<sup>a</sup> The Ministry of Education, Iraq. <sup>b</sup> Faculty of Science, University of Gezira, Public University in Sudan, Sudan.

# Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

#### Article Information

DOI: https://doi.org/10.9734/cjast/2024/v43i124471

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128455

**Original Research Article** 

Received: 15/10/2024 Accepted: 17/12/2024 Published: 28/12/2024

# ABSTRACT

Biliary tree infections arise mainly from stasis associated with acute cholecystitis, choledocholithiasis and acalculous cholecystitis, with appropriate drainage and an effective antibiotic regimen being the mainstay of treatment. The aim of this study was to characterize the bacterial infection of bile cultures in Iraqi patient who admitted to hospital.

Methods. An observational, descriptive, cross-sectional and retrospective study was conducted on 78 patients undergoing cholecystectomy, endoscopic retrograde cholangiopancreatography and cholecystostomy at AL-Anbar Teaching Hospital, Anbar health directorate and Arrazzi Hospital, who

**Cite as:** Hussein, Mohammed Ali, Khalid Elhussein M. Idris, and Mutaman Ali A. Kehail. 2024. "The Most Common Bacterial Infection Among Iraqi Patients Suffering from Cholecystitis in AL-Anbar Province: A Cross-Sectional Observational Study". Current Journal of Applied Science and Technology 43 (12):199-204. https://doi.org/10.9734/cjast/2024/v43i124471.

<sup>\*</sup>Corresponding author: E-mail: moh.bio800@gmail.com;

underwent bile culture the study beginning from March 2022 to May 2023. Participants ranged in age from 10 to 85 years and included both males and females.

Results. 78 patients were included, of whom 70.51 % were women and the average age was  $(63.1 \pm 1.6)$  while men recorded as 29.49% and the average age was  $(60.46 \pm 2.3)$ . The most frequently isolated germ was E. coli (51.2%). The most commonly used antibiotic was ampicillin plus sulbactam (44.6%) followed by piperacillin tazobactam (40.3%).

Conclusion. Escherichia coli is the germ most frequently isolated in biliary tract infections. There is no clarity regarding the use of antimicrobials prophylactically in this pathology, so it is convenient to generate protocols for taking samples and bile cultures in this population, in order to establish the need for the use of antibiotics and to know the bacterial resistance profiles.

Keywords: Microbial infection; biliary tract infections; cholecystitis; antimicrobial agents.

# 1. INTRODUCTION

Cholecystitis is an inflammation of the gallbladder resulting from obstruction of the cystic duct by an impacted gallstone. Its clinical picture is characterized by constant abdominal pain in the upper right quadrant, nausea, vomiting, and fever, with characteristic findings of pain on palpation in the upper right quadrant. Diagnosis requires clinical and paraclinical aspects. which may include a palpable gallbladder, a positive Murphy sign, and increased inflammatory markers in laboratory results. Imaging studies complete the diagnosis (Wu et al., 2021).

Under normal conditions, bile fluid is aseptic, however, with the coexistence of gallbladder pathology, such as cholecystitis, the composition of bile changes and bacteria grow, which is called bacteriobilia. Infections of the biliary tree arise mainly from stasis associated with acute cholecystitis, choledocholithiasis and acalculous cholecystitis2. The germs that most frequently participate are enterobacteria and anaerobium (Cao et al., 2024). The initial management of all patients includes empirical antimicrobial therapy, analgesia and intravenous fluids, with subsequent surgical intervention by laparoscopic cholecystectomy, which is the definitive treatment and the gold standard for this pathology (Leung et al., 1989).

For the above reasons, it is of permanent interest to hospital institutions to know the prevalence of gallbladder and biliary tract pathology (Yun et al., 2006). The use of antibiotic agents therapeutically and prophylactically in the entity where this study was carried out is high in patients undergoing open or laparoscopic cholecystectomy, which implies a high rate of bacterial infections and microbial resistance (Maldonado-Valderrama et al., 2011).

The objective of this study was to characterize the Iraqi patients that underwent cholecystectomy surgery, as well as to determine the main bacterial microbiota present in biliary fluid cultures and their microbial resistance against antibiotics.

#### 2. MATERIALS AND METHODS

Observational, descriptive, cross-sectional and retrospective study, which was carried out at the AL-Anbar Teaching Hospital, Anbar health directorate and Arrazzi Hospital, Iraq. All patients undergoing cholecystectomy, endoscopic retrograde cholangiopancreatography (ERCP) or cholecystostomy, in whom a bile culture was taken, were included between March 2022 to May 2023, a practice routinely carried out in this institution.

Data collection was carried out by three researchers to ensure homogeneity in obtaining information. Additionally, a form was created with the variables of interest, which was linked to the Google Forms digital tool, and was accessed through the researchers' account permissions. The form was anchored to an Excel® database that was updated in real time as the form was filled out.

The samples of vesicular bile obtained during the trans operative period were sent to the laboratory for inoculation in blood, salt and mannitol agar medium, MacConkey and Chocolate (Gram positive/negative and anaerobic) in Bacteriology of the Central Laboratories; four days later the growth of colonies and their antibiotic sensitivity were verified.

# 3. RESULTS

A total of 78 patients were studied, 55 women (70.51%) and 23 men (29.49%), with an average age of 61.4 years (SD  $\pm$  11.3). The body mass index (BMI) was 27.6 (SD  $\pm$  3.4), where 15 patients (19.23%) had normal weight, 42 cases were overweight (53.8%), 16 cases had type I obesity (20.51%) and five patient had type 2 obesity (6.41%) (Figs. 1 and 2).

The preoperative diagnosis was chronic cholecystitis in 53 cases (67.94%), with scheduled surgery. In 25 cases (32.05%), the reason for surgery was acute cholecystitis

(emergency surgery), all of them resolved by laparoscopic cholecystectomy, there were no conversions or transoperative complications.

24 out of the total samples, 30.76%) did not present bacterial growth (Fig. 3) and The most frequent bacteria was proteus mirabilis in 15 samples (19.23%), Staphyloccos areus. in 14 cases (17.94%) and Aeromonas sobria in 3 samples (3.84%). In two patients (2.56%) Enterococcus. faecalis and Staph.lentus were recorded as bacterial infection in bile contents and four isolates of Staph. warneri (5.12%) was reorded Table 1.



Fig. 1. Number and Percentage of patient suffering from cholecystitis related with gender



Fig. 2. Number of Patient related with Body mass index

Type of Microorganisms	NO	acuto	chronic
	<u>NO.</u>	acule	Chronic
Aeromonas sobria	3	3	0
Citrobacter freundii	1	1	0
Enterobcteriacaea	1	1	0
Enterococcus.faecalis	2	1	1
Leuconostoc.mesenteroide	1	0	1
Pantoea ssp	1	0	1
Proteus mirabilis	15	7	8
Providencia stuartii	1	1	0
Pseduomonas aeruginosa	6	1	5
Raoultella.planticola	1	0	1
Rhizobium radiobacter	1	0	1
Serratia marcescens	1	1	0
Staph.aureus	14	12	2
Staph.lentus	2	0	2
Staph.warneri	4	1	3
No growth	24	13	11

Table 1	The most common bacterial infection in	patient with	cholecystitis relat	ed with type of
	inflamm	ation		



Fig. 3. Microbial resistant against different types of antibiotics

Gram-negative microorganisms were reported to to antimicrobials be sensitive such as carbapenems (imipenem), aminoglycosides (gentamicin), cephalosporins, and also to fluorinated quinolones (ciprofloxacin), and for Enterococci and Streptococci, sensitivity to penicillin, cephalosporins, vancomycin, and clindamycin. According to the antibiogram, most of the cultured strains (90%) were sensitive to ceftriaxone (third-generation cephalosporin), and although ciprofloxacin was used as а prophylactic antibiotic in this series of cases, it was only sensitive in 75% of the cases with bacterial development Fig. 3.

Only one emergency case with development of Proteus mirbailisi presented an increase in leukocytes and fever, so the medication was continued for five days, and the patient was discharged without complications.

#### 4. DISCUSSION

Microbial infection is generally associated with advanced age; in our series, the highest number of cases was found after age 51 and is frequently associated with surgical wound infection. Gender is not statistically related to the culture result (Littlefield & Lenahan, 2019). The incidence of positive cultures was with scheduled surgery. In 25 cases (32.05%) acute and 67.94% of cases in chronic cholecystitis), lower than that some previous studeis (15% to 30% in chronic cholecystitis and 32% to 50% in acute cholecystitis) (Wittenburg, 2010) although there is similarity in the bacteria found most frequently: Proteus mirabilis, Escherichia coli, Streptococcus sp. and Enterobacter, form the coliform group of enteric bacteria (Chen et al., 1998).

Considering that cholecystectomy is a cleancontaminated surgery, antibiotic prophylaxis is justified in patients requiring surgery for acute and chronic cholecystitis, due to the frequency of bacteria cultured in bile.9 The initial antibiotic must be effective against the most common infecting microorganisms, coliforms, so in our hospital setting, and based on the sensitivity of the antibiograms performed, cephalosporins seem to be the best option (Sun et al., 2009).

Cholecystectomy is a common surgical procedure. In the area of open cholecystectomy, antibiotic prophylaxis has shown beneficial effects, but these do not seem to be similar in laparoscopic cholecystectomy. There is insufficient evidence to support or refute the use of antibiotic prophylaxis to reduce surgical site infection and systemic infections in patients at low risk of complications (Date et al., 2008).

# 5. CONCLUSION

The rising trend of bacterial infections with drug resistance in patients with cholecystitis in Iraq presents a significant challenge for healthcare providers. This situation is exacerbated by several factors, including inadequate infection control measures, overuse of antibiotics, and limited access to effective treatments.

# DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

# ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

# CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

# **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

# REFERENCES

- Cao, Q., Goldenberg, M., Aron, M., & Wang, T. (2024). Urinary bladder "melanosis": A case report and review of the literature. *International Journal of Surgical Pathology*, 32(1), 104-108.
- Chen, C. Y., Lu, C. L., Huang, Y. S., Tam, T. N., Chao, Y., Chang, F. Y., & Lee, S. D. (1998). Age is one of the risk factors in developing gallstone disease in Taiwan. *Age and Ageing*, *27*(4), 437-441.
- Date, R. S., Kaushal, M., & Ramesh, A. (2008). A review of the management of gallstone disease and its complications in pregnancy. *The American Journal of Surgery, 196*(4), 599-608.
- Leung, J. W., Sung, J. Y., & Costerton, J. W. (1989). Bacteriological and electron microscopy examination of brown pigment stones. *Journal of Clinical Microbiology*, *27*(5), 915-921.
- Littlefield, A., & Lenahan, C. (2019). Cholelithiasis: Presentation and management. *Journal of Midwifery* & *Women's Health, 64*(3), 289-297.
- Maldonado-Valderrama, J., Wilde, P., Macierzanka, A., & Mackie, A. (2011). The role of bile salts in digestion. *Advances in Colloid and Interface Science, 165*(1), 36-46.
- Sun, H., Tang, H., Jiang, S., Zeng, L., Chen, E. Q., Zhou, T. Y., & Wang, Y. J. (2009). Gender and metabolic differences of gallstone diseases. *World Journal of Gastroenterology: WJG, 15*(15), 1886.
- Wittenburg, H. (2010). Hereditary liver disease: Gallstones. Best Practice & Research Clinical Gastroenterology, 24(5), 747-756.
- Wu, Y., Xu, C. J., & Xu, S. F. (2021). Advances in risk factors for recurrence of common bile duct stones. *International Journal of Medical Sciences*, 18(4), 1067.

#### Yun, H. C., Murray, C. K., Roop, S. A., Hospenthal, D. R., Gourdine, E., & Dooley, D. P. (2006). Bacteria recovered from

patients admitted to a deployed US military hospital in Baghdad, Iraq. *Military Medicine*, 171(9), 821-825.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128455