

International Research Journal of Pure and Applied Chemistry

Volume 26, Issue 1, Page 9-22, 2025; Article no.IRJPAC.128564 ISSN: 2231-3443, NLM ID: 101647669

Groundwater Arsenic Contamination and Human Health Risks: A Case Study from the Tongon Gold Mining Area, Côte d'Ivoire

Mahamadou Traore ^a, Dehoule N'Guessan Fulgence Kouassi ^b, Assy Eudes Yapi ^b and N'guessan Louis Berenger Kouassi ^{c*}

^a Département de Géosciences, UFR Sciences Biologiques, Université Peleforo GON COULIBALY, BP 1328, Korhogo, Côte d'Ivoire.

^b Département de Géologie Marine et sédimentologie, UFR STRM, Université Félix HOUPHOUËT-BOIGNY, 22 BP 582 Abidjan 22, Côte d'Ivoire. ^c Département de Mathématiques, Physique et Chimie, UFR Sciences Biologiques, Université Peleforo GON COULIBALY, BP 1328, Korhogo, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. MT, AEY and N'gLBK conceptualized, did the data curation, did formal analysis, did the investigation, did the methodology, did the validation, wrote original draft, wrote, reviewed and edited the manuscript. DN'GFK did data curation, did formal analysis, did the investigation, did the methodology, did the validation, wrote original draft, wrote, reviewed and edited the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/irjpac/2025/v26i1894

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128564

> Received: 22/10/2024 Accepted: 25/12/2024 Published: 10/01/2025

Original Research Article

*Corresponding author: E-mail: kberenger2015@gmail.com;

Cite as: Traore, Mahamadou, Dehoule N'Guessan Fulgence Kouassi, Assy Eudes Yapi, and N'guessan Louis Berenger Kouassi. 2025. "Groundwater Arsenic Contamination and Human Health Risks: A Case Study from the Tongon Gold Mining Area, Côte d'Ivoire". International Research Journal of Pure and Applied Chemistry 26 (1):9-22. https://doi.org/10.9734/irjpac/2025/v26i1894.

ABSTRACT

Human exposure to arsenic (As) is a worldwide concern due to its toxicity, persistence and nonbiodegradability. This exposure can cause cancer and neurological disorders. This study focused on assessing human health risks from arsenic in groundwater near Tongon industrial mining area. The approach consisted firstly to determine total As concentrations in groundwater. Next, groundwater quality was ascertained through water quality index. Finally, the health risks due to As were assessed using carcinogenic and non-carcinogenic risks indices. The sampling was carried out in five wells from May 2023 to July 2023. The results showed that total concentrations of As in the majority wells (14.46 - $32.04 \mu g/L$) are above its WHO guideline value (10 $\mu g/L$). Water quality index values showed good water quality for the wells P2 and P5. While, waters from wells P1, P3 and P4 are poor quality. The non-carcinogenic indices indicated that children are the most exposed to adverse effects through the consumption of wells water around Tongon mining area. The total carcinogenic risk revealated that children and adults can develop cancer. Therefore, it is important to treat wells waters before using for domestic purposes.

Keywords: Groundwater; gold mining; arsenic; human health risk.

1. INTRODUCTION

Poverty reduction through the creation of jobs via mining activities has led to the massive population settling in the vicinity of these mining areas (Taiwo and Awomeso 2017, Traore et al. 2021, Traore et al. 2022). In most of the villages close to these mining areas, groundwater is used by population for various purposes. The quality of these groundwater can be impacted by releasing chemical pollutants via anthropogenic activities such as mining and agricultural activities (Solgi and Jalili 2021, Torrance et al. 2021, Traore et al. 2021, Obodai et al. 2023). Among these pollutants, trace metals and metalloids are a world-wide concern due to their toxicity and bioaccumulation. Hence, trace metals and metalloids monitoring in water is necessary.

Arsenic ingestion by human can cause kidney, lungs, bladers, skin and liver cancers. In addition, respiratory issues, cardiovascular disorders and diabetes are associated to arsenic toxic effects (Kaur et al. 2024, Zhao et al. 2023). Furthermore, Human prolonged exposure via ingestion and dermal contact to arseniccontaminated water can impact negatively their health. (Patel et al. 2023; Ogarekpe et al. 2023, Qaiser et al. 2023;). Therefore, it is important to assess the contamination level of arsenic in groundwater.

In many countries such as Iran (Solgi and Jalili 2021), Turkey (Gemici et al. 2008), Mexico (Esteller et al. 2015), China (Li et al. 2021), and South Africa (Mudzielwana et al. 2020) high arsenic concentrations were found in groundwater around mining areas. The ingestion of these enriched arsenic water by Humans can lead to severe contamination. For example, high concentrations of arsenic were obtained in the urine of residents around Tarkwa mining area $(34 - 700 \ \mu g/L)$ in Ghana (Asante et al. 2007), Mongolia mining area (23.89 - 36.93 $\mu g/L)$ in Korea (Surenbaatar et al. 2021), Akyem mining area (33.95 - 104.97 $\mu g/L)$ in Ghana (Adu-Poku et al. 2019).

Côte d'Ivoire is a country located in west Africa. In this country, during the last two decades, minina activities are grown exponentially. Studies done on arsenic around the mining areas in Côte d'Ivoire were generally focused on soil, sediments and surface waters (Kinimo et al. 2018, Kone et al. 2019, Akpo et al. 2020, Kouakou et al. 2020, Kagbagnan et al. 2023, Kone and Kouakou 2023, N'goran et al. 2023). These studies reported high concentrations of arsenic in soils, sediments and water. For example, Kinimo et al (2018), and N'goran et al (2023) reported that sediments around mining areas in southeastern, and northern Cote d'Ivoire. respectively, were strongly contaminated by arsenic. Soils near the Tongon mining area in the northern Côte d'Ivoire, were extremely contaminated by arsenic (Kone and Kouakou 2023). In addition, Kone et al (2019) reported that surface water in the vicinity of the Tongon mining area were severely contaminated by arsenic. However, data related to Human exposure to arsenic through ingestion or dermal contact of groundwater around mining areas in Côte d'Ivoire are limited. The objectives of the current work were to: (i) determine arsenic concentration in groundwater; (ii) assess the

groundwater quality using water quality index, and (iii) estimate the health risks via noncarcinogenic risk and carcinogenic risk indexes. Data from this study will help in the implantation of groundwater management strategies around mining areas.

2. MATERIAL AND METHODS

2.1 Study Area, Sampling and Laboratory Analysis

The Tongon industrial mining area is located in the northern Côte d'Ivoire, West Africa (Fig. 1). The geology of this zone is dominated by feral soils which are derived from felsic or parent rocks such us granitic, gneiss, phyllite and schist (N'goran et al. 2022). This industrial mine is located from 2.5 km to the Tongon village. The study area is drained by Bandama River which is the largest drainage basins in the country, and the groundwater follows from topographically high areas, in the south-west direction (Sako et al. 2018) The wells P1, P2, P3 and P4 are located around the village of Tongon. While, P5 is located in the village. The agricutural activities namely cotton, cashew, yam, corn, and vegetables crops are carried out around the wells P1 and P3. The well P3 is located in an artisanal gold mining zone. The well P4 is closest to the Tongon industrial gold mining site. Also, the wood coal production and cattle rearing activities are carried out in the vicinity of the well P4. The sampling was done from May 2023 to July 2023. After water collection according the protocol described by Traore et al (2021), samples were acidified with 1 mL of nitric acid HNO₃ (65% suprapur, E.Merck, Germany). Before acidification, the pH of water and the total dissolved solids (TDS) were determined in situ using a multiparameter HI.9828. The major cations (Na⁺, Mg²⁺ and Ca²⁺) and arsenic (As) concentrations were determined through an Inductively Couple Plasma Mass Spectrometer (ICP-MS). The analysis was done in triplicate with an error less than 5%. The map from the study area was done using QGIS software.

2.2 Groundwater Quality and Human Health Risks Assessment

2.2.1 Groundwater quality assessment

The water quality index (WQI) proposed by Sener et al (2017) and Traore et al (2021) was used to assess the quality of the groundwater. WQI is calculated using the Eq. (1):

$$WQI = \sum (W_i \times \frac{C_i}{S_i} \times 100)$$
 (1)

$$W_i = \frac{w_i}{\sum w_i}$$
(2)

With:

 C_i : the determined value of ith parameter S_i : WHO standard value W_i : the relative weight to the element i W_i : the attributed weight to the element i

The WQI < 50 indicates excellent quality. The water quality is good when $50 \le WQI < 100$. The quality is poor for $100 \le WQI < 200$. When $200 \le WQI < 300$, the water quality is very poor. The WQI ≥ 300 shows that the water is undrinkable (Şener et al. 2017, Traoré et al. 2021). The Table 1 gives the S_i, W_i and w_i values.

2.2.2 Human health risks assessment

The hazard index (HI) expressed by the Eq. (3) is used to estimate the total non-carcinogenic risk (Zhang et al. 2017).

$$HI = HQ_{ing} + HQ_{der}$$
(3)

With:

 HQ_{ing} : hazard quotient via ingestion HQ_{der} : hazard quotient through dermal contact When HI < 1, human is exposed to low adverse effects; HI > 1 shows adverse effects (Zhang et al. 2017, Traore et al. 2022).

$$HQ_{ing} = \frac{C_i \times IR \times EF \times ED}{BW \times AT_{nc} \times RfD_{ing}}$$
(4)

$$HQ_{der} = \frac{C_i \times SA \times K_P \times ET \times EF \times ED \times CF}{BW \times AT_{nc} \times RfD_{der}}$$
(5)

$$RfD_{der} = RfD_{ing} \times ABS_{g}$$
 (6)

RfD_{ing} and RfD_{der} are reference dose via ingestion and dermal contact, respectively.

The total carcinogenic risk (TCR) was computed using the Eq. (7) (Zhang et al. 2017)

$$TCR = CR_{ing} + CR_{der}$$
(7)

CR_{der} and CR_{ing} represent the carcinogenic risks via dermal contact and ingestion, respectively. CR_{der} and CR_{ing} are computed as follows:



Fig. 1. Study area and sampling stations

Parameters	WHO standard (2017)	Attributed weight (w _i)	Relative weight (W _i)
рН	6.5-8.5	4	0.2105
Total dissolved solid (TDS)	1000 (mg/L)	4	0.2105
Ca ²⁺	300 (mg/L)	2	0.1053
Mg ²⁺	30 (mg/L)	2	0.1053
Na⁺	200 (mg/L)	2	0.1053
As	0.01 (mg/L)	5	0.2632
		$\sum w_i = 19$	$\sum W_{i} = 1$

Table 1. WHO standard values, and, relative and attributed weights values

Table 2. Exposure assessment parameters

Parameters	Meaning	Unit	Value		Reference
			Adult	Children	
IR	Ingestion rate	L/day	2.2	1	USEPA (2013)
ED	Exposure duration	years	30	10	USEPA (2013)
EF	Exposure frequency	days/year	365	365	USEPA (2013)
BW	Average body weight	kg	70	15	USEPA (2013)
SA	Exposure skin area	cm ²	5700	2800	USEPA (2013)
KP	Permeability coefficient	Cm/h	0.001	0.001	USEPA (2013)
ET	Exosure time	h/day	0.58	1	USEPA (2013)
AT_{nc}	Average time for non- carcinogenic	day	ED×365	ED×365	Coulibaly et al., (2022)
AT _{ca}	Average time for carcinogenic	day	LE×365	LE×365	Coulibaly et al., (2022)
CF	Conversion factor		0.001	0.001	USEPA (2013)
RfD _{ing}	Reference dose of arsenic through ingestion	mg/kg/day	0.3	0.3	USEPA (2013)
SFing	Slop factor of arsenic through ingestion	mg/kg/day	1.5	1.5	USEPA (2013)
ABSg	Gastrointestinal absorption factor		1	1	USEPA (2013)

Life expectancy in Ivory Coast is 58 years (LE = 58 years) (Word Bank, 2019)

$$CR_{ing} = \frac{C_i \times IR \times EF \times ED \times SF_{ing}}{BW \times AT_{ca}}$$
(8)

$$CR_{der} = \frac{C_i \times SA \times K_P \times ET \times EF \times ED \times CF \times SF_{der}}{BW \times AT_{ca}}$$
(9)

$$SF_{der} = \frac{SF_{ing}}{ABS_g}$$
(10)

The carcinogenic risk is acceptable when for TCR range of $10^{-6} - 10^{-4}$. When TCR $\leq 10^{-6}$, the risk is no significant. While, TCR $\geq 10^{-4}$ indicates the possibility for human to develop cancer (Zhang et al., 2017, Traore et al. 2022).

The exposure parameters used to compute the carcinogenic and non-carcinogenic risks indexes are summarized in the Table 2.

3. RESULTS AND DISCUSSION

3.1 Physicochemical Parameters of Groundwater

The pH values and the concentrations of TDS, Na⁺, Mg²⁺ and Ca²⁺ are shown in the Fig. 2. The mean values of pH (Fig. 2e) were 7.63 \pm 0.03, 7.88 \pm 0.29, 7.88 \pm 0.15, 7.06 \pm 0.36, and 7.60 \pm 0.74, in stations P1, P2, P2, P4, and P5, respectively. These values are in the WHO guideline value (6.5 – 8.5), indicating that these waters could be used for domestic purposes (Traoré et al. 2021). The TDS values (Fig. 2d) were below the permissible value (1000 mg/L) of

WHO. Indeed, TDS average values varied between 23.33 ± 5.77 and 213.33 ± 11.54 mg/L. Sodium (Na⁺) concentrations (Fig. 2a) varying from 2.28 mg/L to 19.79 mg/L did not exceed the limit of 200mg/L, showing low decomposition of igneous rocks (Sunitha and Reddy 2019). These finding corroborate those of Traore et al (2021) in the groundwater near the artisanal mining areas in the central-western Côte d'Ivoire. Magnesium (Mg²⁺) concentrations (Fig. 2b) in the stations P2 (16.00 ± 17.62 mg/L), P4 (14.26 ± 15.86 mg/L), and P5 (7.68 ± 6.90 mg/L) were below 30 mg/L (WHO guideline value). Whereas, those in the stations P1 (57.46 ± 88.58 mg/L) and P3 (70.87 ± 113.50 mg/L) exceeded 30 mg/L. According to Singh et al (2012) and Sunitha et al (2019), the decomposition of the ferromagnesia, minerals, the metamorphic rock and the magnesium carbonate in sedimentary rock may release an important amount of Mg²⁺ in the environment. Hence these phenomenon could explained the high concentrations Mg²⁺ obtained in stations P1 and P3. Calcium (Ca2+) concentrations (Fig. 2c) varying from 11.23 ± 10.74 mg/L to 59.34 ± 82.93 mg/L in all the sampling stations were below the WHO limit value of 300 mg/L, indicating low dissolution of calcite and dolomite (Sener et al. 2017).

3.2 Arsenic Distribution in Groundwater

The mean total concentrations of As were 32.04 ± 54.63 µg/L, 14.46 ± 21.67 µg/L, 15.93±20.34 µg/L, 26.92±44.90 µg/L, and 9.75±14.34 µg/L in the groundwater from the stations P1, P2, P3, P4, and P5, respectively (Fig. 2f). The total As concentrations in wells P1, P2, P3 and P4 were higher than the WHO standard (10 μ g/L) (2017), indicating that As may impact negatively human health. In addition, As high concentrations obtained in the wells can be explained by the high infiltration of water and leachates from Tongon mine due to rainfall (Abiye and Bhattacharya 2019, Shams et al. 2020). During the study period, As concentrations did not varied significantly (p < 0.05) among the wells. However, station P1 has registered the highest concentrations. Data (9.75 - 32.04 µg/L) from this study were compared to those from the literature. As total concentrations from the present study were lower than those obtained by Mudzielwana et al (2020) (32.21µg/L) in South Africa, and by Gemici et al (2008) (33 -911 µg/L) in Turkey. Whereas, data found in India by Reddy and Sunitha (2023) (10.3 µg/L), by Sahoo and Khaoash (2020) (0.3 µg/L) and by Singh et al (2018) (2.4 µg/L), in China by Qiao et al

(2020) (0.0027 – 0.003 μ g/L) and by Pan et al (2022) (1.3 μ g/L), in Iran by Solgi and Jalili, (2021) (30.16 μ g/L), and in South Africa by Erdogan et al (2020) (10 – 20 μ g/L) were lower than ours.

3.3 Groundwater Quality and Human Health Risks Assessment

As reported on Fig. 3, the mean values of the water quality index WQI were 161.20 ± 221.93 , 86.13 ± 64.98 , 117.08 ± 62.88 , 122.25 ± 143.13 , and 65.76 ± 45.61 in stations P1, P2, P3, P4 and P5, respectively, thereby suggesting poor quality of waters from the wells P1, P3, and P4. Whereas, waters from wells P2 and P5 qualities are good.

Table 3 indicates the computed values of noncarcinogenic risk (HQing, HQder, HI) and carcinogenic risk (CRing, CR der, TCR) indexes. The HI values for adults and children were ranged from 0.44 to 1.44, and from 2.17 to 7.14, respectively. All HI values for children were above 1. For adults, the HI value obtained at stations P1 (1.44) and P4 (1.21) were higher than 1. These results showed potential adverse effects for children, and adults. The results also revealed that the values of HQing (above 1 for children at all stations, and for adults at stations P1 and P4) were higher than those of HQder, showing that arsenic induced adverse effects through ingestion. In addition, children are the most exposed to adverse effects due to their low body weight compared to that for adults, and to their rapid growing (Slogi and Jalili 2021). Indeed, the adults's HI values were lower than those of children. The mean values for CRing varied between 2.38×10^{-4} and 7.81×10^{-4} for adults, and, between 1.68×10⁻⁴ and 5.52×10⁻⁴ for children. Those for CRder ranged from 6×10⁻⁷ to 10⁻⁶ for adults, and from 4.7×10^{-to 7}10⁻⁶ for children. The values of TCR, varied from 2.38×10^{-4} and 7.82×10^{-4} for adults, and from 1.69×10^{-4} and 5.52×10^{-4} for children. The values of CRing and TCR both for children and adults were above 10⁻⁴, suggesting that children and adults can develop cancer.

3.4 Sources of Arsenic in Groundwater

To know the possible sources of As in the groundwater, principal component analysis was performed. The results are shown in Table 4. Three PC loadings accounted for 44.25 % (PC1), 35.99 % (PC2) and 12.28 % (PC3) were selected. The major cations Na⁺, Mg²⁺, and Ca²⁺

			P1	P2	P3	P4	P5
		HQing	1.44 ± 2.45	0.65 ± 0.97	0.72 ± 0.91	1.21 ± 2.02	0.44 ± 0.65
Non-	Adult	HQder	2.16×10 ⁻³ ± 3.69×10 ⁻³	9.75×10 ⁻⁴ ± 1.46×10 ⁻³	1.08×10 ⁻³ ± 1.37×10 ⁻³	1.82×10 ⁻³ ± 3.03×10 ⁻³	6.58×10 ⁻⁴ ± 9.7×10 ⁻⁴
carcinogenic		HI	1.44 ± 2.46	0.65 ± 0.97	0.72 ± 0.91	1.21 ± 2.02	0.44 ± 0.65
-		HQing	7.12 ± 12.14	3.21 ± 4.83	3.54 ± 4.52	5.98 ± 9.98	2.17 ± 3.20
	Children	HQder	1.99×10 ⁻² ± 3.4×10 ⁻²	9.00×10 ⁻³ ±1.35×10 ⁻²	9.91×10 ⁻³ ± 1.27×10 ⁻²	1.67×10 ⁻² ± 2.79×10 ⁻²	6.1×10 ⁻³ ±8.9×10 ⁻³
		HI	7.14 ± 12.17	3.22 ± 4.83	3.55 ± 4.53	6.00 ± 10.01	2.17 ± 3.20
		CRing	7.81×10 ⁻³ ±1.33×10 ⁻³	3.53×10 ⁻⁴ ± 5.3 ×10 ⁻⁴	3.88×10 ⁻⁴ ± 4.95×10 ⁻⁴	6.56×10 ⁻⁴ ± 1.1×10 ⁻⁴	2.38×10 ⁻⁴ ± 3.5×10 ⁻⁴
Carcinogenic	Adult	CRder	1.20×10 ⁻⁶ ± 2.00×10 ⁻⁶	$5.00 \times 10^{-7} \pm 7.94 \times 10^{-7}$	5.83×10 ⁻⁷ ± 7.45×10 ⁻⁷	9.86×10 ⁻⁷ ± 1.64×10 ⁻⁶	3.57×10 ⁻⁷ ± 5.26×10 ⁻⁷
-		TCR	7.81×10 ⁻³ ±1.33×10 ⁻³	3.53×10 ⁻⁴ ± 5.3 ×10 ⁻⁴	3.88×10 ⁻⁴ ± 4.95×10 ⁻⁴	6.57×10 ⁻⁴ ± 1.1×10 ⁻⁴	2.38×10 ⁻⁴ ± 3.5×10 ⁻⁴
		CRing	5.52×10 ⁻⁴ ± 9.41×10 ⁻⁴	2.49×10 ⁻⁴ ± 3.73×10 ⁻⁴	2.75×10 ⁻⁴ ± 3.5×10 ⁻⁴	4.64×10 ⁻⁴ ± 7.74×10 ⁻⁴	1.68×10 ⁻⁴ ± 2.48×10 ⁻⁴
	Children	CRder	2.00×10 ⁻⁶ ± 2.64 ×10 ⁻⁶	6.97×10 ⁻⁷ ± 1.04×10 ⁻⁶	7.69×10 ⁻⁷ ± 9.81×10 ⁻⁷	1.29×10 ⁻⁶ ± 2.16×10 ⁻⁶	4.70×10 ⁻⁷ ± 6.97×10 ⁻⁷
		TCR	5.54×10 ⁻⁴ ± 9.41×10 ⁻⁴	2.49×10 ⁻⁴ ± 3.73×10 ⁻⁴	2.75 ×10 ⁻⁴ ± 3.5×10 ⁻⁴	4.65×10 ⁻⁴ ± 7.74×10 ⁻⁴	$1.68 \times 10^{-4} \pm 2.48 \times 10^{-4}$

Table 3. Values of non-carcinogenic risk and carcinogenic risk indices



Traore et al.; Int. Res. J. Pure Appl. Chem., vol. 26, no. 1, pp. 9-22, 2025; Article no.IRJPAC.128564

Fig. 2. Spatial variations of sodium (a), magnesium (b), calcium (c), total dissolved solids (d) and arsenic (f) concentrations, and (e) spatial variations of pH

Wells

P1 P2 P3 P4 P5

Wells

P1 P2 P3 P4 P5

P1 P2 P3 P4 P5

Wells



Fig. 3. Spatial variation of water quality index (WQI)

were bound strongly on PC1 with loading values of 0.96, 0.86, and 0.96, respectively. This implies that these elements could have a common origin. According to Rakotondrabe et al. (2018), low concentrations of major cations compared to their WHO guidelines values are due to the natural processes such us silicate rocks and carbonate minerals dissolution. In this study, Na⁺, Mg²⁺, and Ca²⁺ concentrations are below to their WHO permissible levels. Therefore, these are naturally released in elements the groundwater. Hence, PC1 can be interpreted as a lithological contribution to the groundwater around Tongon mining area chemistry. The parameters pH and TDS, as well as arsenic contribute to the PC2. The loading values were 0.94 for pH, 0.89 for TDS and (-0.68) for As. It has been reported that, when the solubility of trace metal decresed under basic pH, the trace metal is measured in low concentration (Brindha et al., 2020). In the present study, the majority groundwaters pH are alkaline and the As total concentration are higher the WHO permissible level. Therefore, the pH not influence As distribution in the groundwater. That may explain the negative contribution of As on the PC 2. On the PC 3, only As was bound with loading value of 0.69, indicating that As in groundwater may derived from anthropogenic activities. This can be explained by the lack loading of Na⁺ and Ca²⁺ on the PC3. Indeed, an exchange of Na⁺ and

Ca²⁺ lead to high As concentration in aroundwater. This exchange occurs when the ratio of Na⁺/Ca²⁺ is above to 1 (Sako et al., 2016). The Na⁺/Ca²⁺ ratios in the all groundwater around the tongon mining area varied between 0.08 and 0.31 (Fig. 4). Therefore, there is no exchange of Na⁺ for Ca²⁺. Hence the high concentrations of As can not be explained by the exchange of Na⁺ for Ca²⁺. Na⁺ and Ca²⁺ not play an important role in the distribution of As in the groundwater. In the study area, industrial and artisanal gold mining activities are well developed. In addition, population use pesticides and phosphate fertilizers to improve the production of the cotton, cashew, yam, corn, and vegetables crops. These agrochemicals release an important amount of As in the environment (Javasumana et al. 2015, Patel et al. 2023). Therefore, the agricultural and gold minining activities can be potential sources of As in the groundwater. The results of pearson correlation analysis (Table 5) indicates that Mg²⁺ and Ca²⁺ are correlated (r = 0.72), Na⁺ is correlated with Ca^{2+} (r = 0.97) and with Mg²⁺ (r = 0.71). In addition, pH and TDS are correlated (r = 0.84). Whereas, no correlation was found between As and the major cations (Mg²⁺, Ca²⁺, and Na⁺), and between the parameters (pH and TDS). These results are in accordance with those of PCA analysis.



Wells

Fig. 4. Sapatial variation of the ratio Na/Ca

	PC 1	PC 2	PC 3	
As	0.17	-0 .68	0.69	
Na	0.96	0.09	-0.22	
Mg	0.86	-0.03	0.23	
Ca	0.96	-0.11	-0.16	
pH	0.002	0.94	0.26	
TDS	0.18	0.89	0.26	
Eigen value	2.65	2.16	0.74	
Total variance (%)	44.25	35.99	12.28	
Cumulative variance (%)	44.25	80.24	92.52	

Table 4. Principal component analysis (PCA) results

Table 5. Correlation coefficients (p<0.05; n = 15) of major cations (Na⁺, Mg²⁺, Ca²⁺), arsenic (As), pH and total dissolved solids (TDS)

	As	Na	Mg	Са	рН	TDS	
As	1.00						
Na	-0.02	1.00					
Mg	0.24	0.71	1.00				
Ca	0.16	0.97	0.72	1.00			
pН	-0.49	-0.0004	0.11	-0.17	1.00		
TDS	-0.33	0.24	0.05	0.09	0.84	1.00	

4. CONCLUSION

This study focused on assessing health risks from arsenic in groundwater near Tongon industrial mining area. The results indicated that pH values are in the WHO guideline value (6.5 -Sodium (Na⁺) and calcium (Ca²⁺) 8.5). concentrations were lower than thier WHO guideline values (200 mg/L for Na⁺ and 300 mg/L for Ca²⁺). Magnesium (Mg²⁺) concentrations exceeded its guideline value of 30 mg/L in the wells P1 and P3. The results also showed that As concentrations did not varied significantly (p < 0.05) among the wells. The total As concentrations in the wells P1 (32.04 \pm 54.63 µg/L), P2 (14.46 ± 21.67 µg/L), P3 (15.93±20.34 μ g/L), and P4 (26.92 ± 44.90 μ g/L) were above the WHO standard (10 µg/L). Water quality index values showed good water quality for the wells P2 and P5. While, waters from wells P1, P3 and P4 are poor quality. The results of noncarcinogenic risk indices showed potential adverse effects for children through the water from all the wells, and adults through the water from the wells P1 and P4. The total carcinogenic risk suggested that children and adults can develop cancer. Complementary studies including arsenic concentrations assessment in the urine, blood and hair must be done to better understand the deleterious effects on the residents around the Tongon mining area.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The authors express their gratitude to the President of Peleforo GON COULIBALY University, Korhogo for facilitating the experiments in the laboratory.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abiye, T. A., & Bhattacharya, P. (2019). Arsenic concentration in groundwater: Archetypal study from South Africa. *Groundwater for Sustainable Development, 9*, 100246. https://doi.org/10.1016/j.gsd.2019.100246
- Adu-Poku, B., Asiedu, N., Akoto, O., & Ataki, J. (2019). Modelling the distribution of arsenic and mercury in urine using chemometric tools. *Cogent Chemistry*, *5*, 1586064. https://doi.org/10.1080/23312009.2019.158 6064

- Akpo, K. S., Koné, T., Coulibaly, S. L., Kamagate, M., & Coulibaly, L. (2020). Diagnostics of environmental risks and mapping of surface water sensitivity due to metal contamination from artisanal gold mining in Côte d'Ivoire: Case of Angovia, Kokumbo, Hire, and Agbaou. *Environmental Protection Science*, 8, 47– 64. https://doi.org/10.4236/gep.2020.82004
- Asante, K. A., Agusa, T. A., Subramanian, A., Ansa-Asare, O. D., Biney, C. A., & Tanabe, S. (2007). Contamination status of arsenic and other trace elements in drinking water and residents from Tarkwa, a historic mining township in Ghana. *Chemosphere*, *66*, 1513–1522. https://doi.org/10.1016/j.chemosphere.200 6.08.022
- Brindha, K., Rajib, P., Walter, J., Tan, M. L., & Singh, M. K. (2020). Trace metals contamination in aroundwater and implications on human health: Comprehensive assessment usina hydrogeochemical geostatistical and methods. Environmental Geochemistry Health. 3819-3839. and 42, https://doi.org/10.1007/s10653-020-00637q
- Coulibaly, M., Kouassi, N. L. B., N'goran, K. P. D. A., Diabate, D., & Trokourey, A. (2022). Distribution, ecological, and health risks of arsenic in sediment from the mixing zone of the Comoé River and the Ébrié Lagoon, Côte d'Ivoire, West Africa. *American Journal of Applied Chemistry*, *10*, 89–96. https://doi.org/10.11648/j.ajac.20221004.1 3
- Erdogan, I. G., Fosso-Kankeu, E., Ntwampe, S. K. O., Waanders, F., & Hoth, N. (2020). Seasonal variation of hydrochemical characteristics of open-pit groundwater near a closed metalliferous mine in O'Kiep, Namaqualand Region, South Africa. *Environmental Earth Sciences, 79*, 119. https://doi.org/10.1007/s12665-020-8863-2
- Esteller, M. V., Dominguez-Mariani, E., Garrido, S. E., & Avilés, M. (2015). Groundwater pollution by arsenic and other toxic elements in an abandoned silver mine, Mexico. *Environmental Earth Sciences*, 74, 2893–2906.

https://doi.org/10.1007/s12665-015-4315-9

Gemici, U., Tarcan, G., Helvacı, C., & Somay, A. M. (2008). High arsenic and boron concentrations in groundwaters related to mining activity in the Bigadic borate deposits (Western Turkey). *Applied* *Geochemistry,* 23, 2462–2476. https://doi.org/10.1016/j.apgeochem.2008. 02.013

- Jayasumana, C., Fonseka, S., Fernando, A., K., Amarasinghe, Μ., Jayalath, Siribaddana, Gunatilake, S., & S., Paranagama, Ρ. (2015). Phosphate fertilizer is a main source of arsenic in areas affected with chronic kidney disease unknown etiology in Sri Lanka. of SpringerPlus. 4. 90. http://dx.doi.org/10.1186/s40064-015-0868-z
- Kagbagnan, K., Adjoumani, R. K., Adou, K. E., Ehouman, A. D., Yao, B., & Adouby, K. (2023). Evaluation of metal contamination of sediments around an industrial gold mine in Côte d'Ivoire: The example of arsenic. *Journal of Applied Science and Environmental Management,* 27, 1627– 1632.
- Kaur, R., Garkal, A., Sarode, L., Bangar, P., Mehta, T., Singh, D. P., & Rawal, R. (2024). Understanding arsenic toxicity: Implications for environmental exposure and human health. *Journal of Hazardous Materials Letters*, *5*, 100090. https://doi.org/10.1016/j.hazl.2023.100090
- Kinimo, K. C., Yao, K. M., Marcotte, S., Kouassi, N. L. B., & Trokourey, A. (2018).
 Distribution trends and ecological risks of arsenic and trace metals in wetland sediments around gold mining activities in central-southern and southeastern Côte d'Ivoire. *Journal of Geochemical Exploration, 190, 265–280.* https://doi.org/10.1016/j.gexplo.2018.03.01 3
- Kone, K., & Kouakou, A. R. (2023). Assessment of soil contamination and human health risk around an industrial gold mine in Côte d'Ivoire: The case of arsenic. *Open Journal* of Soil Science, 13, 329–339. https://doi.org/10.4236/ojss.2023.137014
- Kone, K., Kouakou, A. R., Kouassi, K. E., & Kopoin, A. (2019). Heavy metal pollution index of surface water and groundwater around Tongon mine (Côte d'Ivoire). *International Journal of Environmental Monitoring and Analysis*, 7, 103–111. https://doi.org/10.11648/j.ijema.20190705. 12
- Kouakou, Y. K. N., Trazié, J. G. B., Coulibaly, J.
 L. K., & Ahou Clarisse Koffi, A. A. (2020).
 Assessment of heavy metals on soils around Gbetogo mine at Séguéla, North-West of Côte d'Ivoire. *International Journal*

of Innovative and Applied Studies, 31, 176–188.

- Mudzielwana, R., Gitari, M. W., Akinyemi, S. A., Talabi, A. O., & Ndungu, P. (2020). Hydrogeochemical characteristics of arsenic-rich groundwater in Greater Giyani Municipality, Limpopo Province, South Africa. *Groundwater for Sustainable Development, 10, 100336.* https://doi.org/10.1016/j.gsd.2020.100336
- N'goran, K. P. D. A., Diabate, D., Kouassi, N. L. B., Yao, K. M., Kinimo, K. C., Ouattara, A. A., & Trokourey, A. (2022). Geochemical speciation and distribution of trace metals in sediments around industrial and artisanal gold mining areas in northern Côte d'Ivoire. *Environmental Earth Sciences, 81*, 333. https://doi.org/10.1007/s12665-022-10447-0
- N'goran, K. P. D. A., Fato, T. P., Kouassi, N. L. B., & Diabate, D. (2023). Ecological and human health risks assessment of arsenic in sediments around gold mining areas in northern Côte d'Ivoire. *Asian Journal of Chemistry Science, 13*, 234–247. https://doi.org/10.9734/AJOCS/2023/v13i6 277
- Obodai, J., Duncan, A. E., Adjei, K. A., & Odai, S. N. (2023). A preliminary investigation of surface and groundwater quality along the upper part of the Ankobra River, impacted by illegal mining activities. *Water, Air, & Soil Pollution, 234*, 162. https://doi.org/10.1007/s11270-023-06170-6
- Ogarekpe, N. M., Nnaji, C. C., Oyebode, O. J., Ekpenyong, M. G., Ofem, O. I., Tenebe, I. T., & Asitok, A. D. (2023). Groundwater quality index and potential human health risk assessment of heavy metals in water: A case study of Calabar metropolis, Nigeria. *Environmental Nanotechnology, Monitoring & Management, 19*, 100780. https://doi.org/10.1016/j.enmm.2023.10078 0
- Pan, H., Zhou, G., Yang, R., Cheng, Z., & Sun, B. (2022). Heavy metals and As in groundwater, surface water, and sediments of Dexing Giant Cu-polymetallic ore cluster, East China. *Water*, 14(3), 352. https://doi.org/10.3390/w14030352
- Patel, B., Gundaliya, R., Desai, B., Shah, M., Shingala, J., Kaul, D., & Kandya, A. (2023). Groundwater arsenic contamination: Impacts on human health and agriculture, ex situ treatment

techniques and alleviation. *Environmental Geochemistry and Health, 45*, 1331–1358. https://doi.org/10.1007/s10653-022-01334-5

- Patel, K. S., Pandey, P. K., Martín-Ramos, P., Corns, W. T., Varol, S., Bhattacharya, P., & Zhu, Y. (2023). A review on arsenic in the environment: Contamination, mobility, sources, and exposure. *RSC Advances*, *13*, 8803–8821. http://dx.doi.org/10.1039/D3RA00789H
- Qaiser, F. U. R., Zhang, F., Pant, R. R., Zeng, C., Khan, N. G., & Wang, G. (2023). Characterization and health risk assessment of arsenic in natural waters of the Indus River Basin, Pakistan. *Science of The Total Environment, 857*, 159408. https://doi.org/10.1016/j.scitotenv.2022.159 408
- Qiao, D., Wang, G., Li, X., Wang, S., & Zhao, Y. (2020). Pollution, sources, and environmental risk assessment of heavy metals in the surface AMD water, sediments, and surface soils around unexploited Rona Cu deposit, Tibet, China. *Chemosphere*, 248, 125988. https://doi.org/10.1016/j.chemosphere.202 0.125988
- Sahoo, S., & Khaoash, S. (2020). Impact assessment of coal mining on groundwater chemistry and its quality from Brajrajnagar coal mining area using indexing models. *Journal of Geochemical Exploration, 215*, 106559–106566. https://doi.org/10.1016/j.gexplo.2020.1065 59
- Sako, A., Bamba, O., & Gordio, A. (2016). Hydrogeochemical processes controlling groundwater quality around Bomboré gold mineralized zone, Central Burkina Faso. *Journal of Geochemical Exploration, 170*, 58–71. http://dx.doi.org/10.1016/j.gexplo.2016.08.

009 Sako, A., Semdé, S., & Wenmenga, U. (2018).

- Geochemical evaluation of soil, surface water, and groundwater around the Tongon gold mining area, northern Côte d'Ivoire, West Africa.
- Şener, S., Şener, E., & Davraz, A. (2017). Evaluation of water quality using the water quality index (WQI) method and GIS in Aksu River (SW-Turkey). Science of The Total Environment, 584–585, 131– 144. https://doi.org/10.1016/j.scitotenv.2017.01.

102

Shams, M., Nezhad, N. T., Dehghan, A., Hossein, A., Paydar, M., Mohammadi, A. A., & Zarei, A. (2020). Heavy metals exposure, carcinogenic and noncarcinogenic human health risks assessment of groundwater around mines in Joghatai, Iran. *International Journal of Environmental Analytical Chemistry*, *102*, 1884–1899.

https://doi.org/10.1080/03067319.2020.174 3835

- L., Singh, U. K., Ramanathan, Α. & Subramanian, V. (2018). Groundwater chemistry and human health risk assessment in the mining region of East Singhbhum, Jharkhand, India. Chemosphere. 501-513. 204. https://doi.org/10.1016/j.chemosphere.201 8.04.060
- Solgi, E., & Jalili, M. (2021). Zoning and human health risk assessment of arsenic and nitrate contamination in groundwater of agricultural areas of the twenty-two villages with geostatistics (Case study: Chahardoli Plain of Qorveh, Kurdistan Province, Iran). *Agricultural Water Management,* 255, 107023.

https://doi.org/10.1016/j.agwat.2021.10702 3

- Sunitha, V., & Reddy, S. Y. (2019). Hydrogeochemical evaluation of groundwater in and around Lakkireddipalli and Ramapuram, Y. S. R District, Andhra Pradesh, India. *HydroResearch, 2*, 85–96. https://doi.org/10.1016/j.hydres.2019.11.00 8
- Surenbaatar, U., Kim, B. G., Seo, J. W., Lim, H. J., Kwon, J. K., Kang, M. K., Altangerel, E., Byambaa, T., Batbaatar, S., Myagmardorj, O., Lee, C. W., & Hong, Y. S. (2021).
 Environmental health survey for children residing near mining areas in South Gobi, Mongolia. *Annals of Occupational and Environmental Medicine*, 33, e10. http://dx.doi.org/10.35371/aoem.2021.33.e 10
- Taiwo, A. M., & Awomeso, J. A. (2017). Assessment of trace metal concentration and health risk of artisanal gold mining activities in Ijeshaland, Osun State Nigeria—Part 1. *Journal of Geochemical Exploration, 177, 1–10.* https://doi.org/10.1016/j.gexplo.2017.01.00 9
- Torrance, K. W., Redwood, S. D., & Cecchi, A. (2021). The impact of artisanal gold mining, ore processing and mineralization

on water quality in Marmato, Colombia. *Environmental Geochemistry and Health,* 43, 4265–4282. https://doi.org/10.1007/s10653-021-00898v

- Traore, Z., Kouassi, N. L. B., Assemian, A. S., Kouassi, K. E., Drogui, P., & Yao, K. B. (2021). Distribution and health risk assessment of trace metals in surface waters and groundwater around artisanal gold mining areas in central-western Côte d'Ivoire, West Africa. *International Journal of Environmental Monitoring and Analysis*, 9, 136–151. https://doi.org/10.11648/j.ijema.20210905. 15
- Traore, Z., Kouassi, N. L. B., Assemian, A. S., Kouassi, K. E., Kopoin, A., & Yao, K. B. (2022). Ecological and human health risks assessment of chromium in soils around artisanal gold mining areas: A case study in central-western Côte d'Ivoire, West Africa. *Chemistry in Africa, 5*, 1863– 1871.

https://doi.org/10.1007/s42250-022-00478-2

- United States Environmental Protection Agency (USEPA). (2013). Regional screening level (RSL) summary table. Available at: http://www.epa.gov/reg3hwmd/risk/human/ rbconcentration_table/Generic_Tables/docs/ master_sl_table_run_NOV2013.pdf. Accessed November 2013.
- World Bank. (2019). The World Bank Group: Average Life Expectancy in Côte d'Ivoire: 2009–2019. https://data.worldbank.org/indicator/SP.DY N.LE00.IN?locations=CI. Accessed 12 July 2021.
- World Health Organization (WHO). (2017). Guidelines for drinking-water quality: Fourth edition incorporating first addendum. World Health Organization. https://apps.who.int/iris/handle/10665/2546 37 (Accessed 28 April 2021).
- Zhang, Y., Chu, C., Li, T., Xu, S., Liu, L., & Ju, M. (2017). A water quality management strategy for regionally protected water through health risk assessment and spatial distribution of heavy metal pollution in 3 marine reserves. *Science of The Total Environment, 599–600, 721–* 731.

https://doi.org/10.1016/j.scitotenv.2017.04. 232 Traore et al.; Int. Res. J. Pure Appl. Chem., vol. 26, no. 1, pp. 9-22, 2025; Article no.IRJPAC.128564

Zhao, C., Du, M., Yang, J., Guo, G., Wang, L., Yan, Y., Li, X., Lei, M., & Chen, T. (2023). Changes in arsenic accumulation and metabolic capacity after environmental

management measures in mining area. Science of The Total Environment, 855, 158652.https://doi.org/10.1016/j.scitotenv.2 022.158652

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128564