

Understanding the Phylogenetics and Evolution of Genus *Schistosoma*- Africa and Asia Stand Point

**Olugbenga Samson Onile^{1,2,4*}, Benson Otarigho²
and Chiaka I. Anumudu³**

¹Biotechnology Unit, Department of Biological Sciences, Elizade University Ilara-Mokin, Ondo State, Nigeria.

²Cell Biology and Genetics Unit, Department of Zoology, University of Ibadan, Nigeria.

³Cellular Parasitology Programme, Department of Zoology, University of Ibadan, Nigeria.

⁴Centre for Biotechnology Resources and Development (CENBRAD), Elizade University Ilara-Mokin, Ondo State, Nigeria.

Authors' contributions

Author OSO designed the study, performed the software analysis, wrote the protocol and wrote the first draft of the manuscript. Author BO help in the design of methodology and also provide some technical assistance during analysis. Author CIA conducted the literature searches, make final review of the analysis and study for correction and provide us with further professional information. All authors read and approved the final manuscript.

Original Research Article

Received 27th September 2013
Accepted 28th November 2013
Published 11th February 2014

ABSTRACT

The evolutionary spread of schistosomes infection was reportedly prominent more in Africa and Asia continents. This study therefore examined the evolutionary trend of this parasite while limiting the investigation to *Schistosoma* species peculiar to this region of the world. The evolutionary history of this species group was inferred using DNA sequences from NCBI Genbank database and Maximum likelihood, Ancestral inference; Neighbor-Joining method analysis was employed in this study. All members of this species complex were AT rich, with *S. mekongi* and *S. malayensis* having the highest AT nucleotide composition. The smallest evolutionary divergence was also observed in *S. curassoni* and *S. bovis*. The finding of this study slightly contradict previous report on ancestral prediction of Schistosomes.

*Corresponding author: Email: onileg@yahoo.com, olugbenga.onile@elizadeuniversity.edu.ng;

Keywords: Phylogenetics; evolution; *Schistosoma*; nucleotide.

1. INTRODUCTION

Human schistosomiasis is endemic in large areas of the sub-tropics. Over 700 million people in 74 countries have been estimated to be exposed to the risk of *Schistosoma* infection and almost 200 million were estimated to be infected in 2003 [1], of which 85% in sub-Saharan Africa [21]. Schistosomiasis is a disease affecting agricultural communities, particularly those dependent upon irrigation to support their agriculture. Evolutionary history of the parasite *Schistosoma* has generated much interest especially with regard to understanding the precise roles that particular species have in causing human disease [2,3,21]. *Schistosoma species* as described by Lawton *et al.*, were categorised into four major groups based on their distribution, host specificity and egg morphology, this include the *S. japonicum* (found throughout eastern and southern Asia), *S. indicum* (inhabits the western and southern regions including India, Thailand and Sri Lanka), *S. mansoni* (Africa) and *S. haematobium* (throughout Africa) group [4]. Phylogenetic and phylogeographic studies can identify historically and evolutionary independent groups of populations and can also identify natural or anthropogenic colonization routes [5]. Analyses of genetic structure use patterns of neutral genetic diversity to reveal how populations are structured across geography, within and among hosts on a local scale. This approach can provide information about pathogen transmission that is difficult to ascertain otherwise [6]. Genetic structure analyses are powerful tools for inferring epidemiological and evolutionary processes in schistosomes. They indicate that transmission foci are structured by watershed boundaries, habitat types, and host species. Recently molecular markers have been used for a variety of genomic-based taxonomic, phylogenetic, population and evolutionary investigations in animal species [7,8]. One of these DNA markers is the Mitochondrial DNA (mtDNA) sequence. Although mtDNA-sequence data have proved valuable in phylogenetic analysis, the selection of the appropriate gene for analysis is important. Among the coding genes in the mitochondrion genome, subunit I of the cytochrome oxidase (COI) gene possesses features suitable for evolutionary studies [9,10]. MtDNA possess a relatively fast mutation rate, which results in significant variation in mtDNA sequences between species providing an ample within species variance useful for phylogenetic investigations [11,12]. The present study, explored by retrieving the cytochrome oxidase subunits I (COI) of mitochondrial DNA of the 16 member of the *Schistosoma species* to infer the evolution and phylogeny of this group while considering Africa and Asia continent perspective.

2. MATERIALS AND METHODS

2.1 Schistosomes Sequence Retrieval

16 nucleotide sequences of individual mtDNA gene sequences of COI and an outgroup *ssrDNA* sequence, of members of the *Schistosoma species* from the GenBank database (www.ncbi.nlm.nih.gov) was retrieved (see Table 1). These were chosen based on their classification as African and Asian oriented schistosoma species [3]. Sequence alignment was performed by using Clustal Omega software (<http://www.ebi.ac.uk/Tools/msa/clustalo>), through elimination of all positional gaps and missing data. The sequences were then trimmed to get their equal lengths for all the species.

2.2 Inferring Evolutionary History

The evolutionary history was inferred using Maximum Likelihood method based on the Tamura-Nei model [13], Neighbor-Joining method [14] and Minimum Evolution method [15]. All the phylogenetic analyses were conducted in MEGA software (<http://www.megasoftware.net>), version 5.0 [16]. Ancestral states were inferred using the Maximum Likelihood method [17], under the General Time Reversible model [17]. All the positions containing gaps or missing data were eliminated (complete deletion) from the dataset prior to analysis. However, MEGA software also provides option to retain all such sites initially and excluding them as necessary in the pair-wise distance estimation (pair-wise deletion option) or to use the partial deletion (site coverage) as a percentage. Evolutionary analyses were conducted in MEGA 5 [16].

2.3 Prediction of Evolutionary Divergence

Evolutionary divergence between and overall Sequences pairs of the *Schistosoma species* were achieved by using the Maximum Composite Likelihood model [17] of the MEGA 5 software.

3. RESULTS AND DISCUSSION

The nucleotide composition of each taxon in the examined family as presented in Fig. 1, shows that all members of this family were all AT rich, which could possibly imply that AT (Adenine-thymine) rich environment in schistosome COI genes revealing long (dA).(dT) tracts that are found preferentially in introns and flanking gene regions, and thus pose an advantage for evolutionary study. *S. mekongi* and *S. malayensis* had the highest AT composition with both having 47.8% thymine(T)/21.8% Adenine(A) and 47.6% thymine (T)/21.8% Adenine (A) respectively, while *S. mansoni* had the least among the family with 41.2% thymine (T) and 27.1% Adenine (A). The nucleotide frequencies are 21.74% (A), 46.78% (T), 21.47% (C) and 10.01% (G). There is more of thymine in the sequences and less of guanine. The estimated evolutionary distance of divergence between sequences of *Schistosoma species* is presented in table 3, with the smallest evolutionary divergence found between *S. curassoni* and *S. bovis* (0.053), which is followed by *S. intercalatum*/*S. curassoni* and *S. intercalatum*/*S. curassoni* (with both groups at evolutionary divergence of 0.064 respectively). The predicted evolutionary divergence is observed to closely related to the outcome of phylogenetic analysis conducted in this study (Figs. 2-5). This result slightly differs from phylogenetic report where this species were duly considered (3). The divergence between other paired species was quite greater. It could be inferred from this result that *S. curassoni* and *S. bovis* are the most closely related taxa. The overall transition/transverse bias is $R=0.823$, where $R=[A^*G^*k_1+T^*C^*k_2]/[(A+G)^*(T+C)]$. The study shows that transverse mutational frequencies were more than that of transitional mutation (Table 2). Transitional mutation occurs more in adenine (16.75), guanine (16.56) and far less occur in cytosine (3.51). The ancestral state was inferred using Maximum Likelihood method [16] and is presented in Fig. 2. It could be incidental that the ancestral origin from which other members of this complex was predicted is *S. mansoni*, being the species with the longest estimated ancestral inference. *S. bovis* and *S. nasale* are the most recently evolved taxa among the *Schistosoma species*, which is followed by *S. mattheei* from where they possibly share similar ancestors. The ancestral node from which *S. indicum* and *S. intercalatum* evolved from was inferred to be *S. malayensis*, which share a very close ancestral linkage with *S. mansoni*. *S. haematobium* and *S. incognitum* also share a close ancestral connection with *S.*

margrebowiei and *S. spindale*. The ancestral prediction of this study shows a migration initiated from Africa via *S. mansoni* through the central and South East Asia (India) and then back to Africa (Figs. 2,3,4,5). To foster our understanding about the ancestral prediction of schistosoma species, an outgroup maximum likelihood phylogenetics analysis was also conducted, using a *Schistosoma* species ssrDNA sequence (Fig. 5), the outcome of this analysis also correspond with other related analysis and thus further buttress the evolutionary migration of Schistosomiasis from Africa to Asia and back to Africa. It could be deduced that lack of records of infection in Africa and cultural challenges while considering the symptoms of this infection may be a challenge in proper report of this infection. Many studies have reported the geographical evolution of this infection [3,19,20] and have reported the Asian ancestry of the schistosomes, and also gave weight to the concept of the reinvasion back from Africa. Evolutionary history inferred using the Maximum likelihood and Neighbour-Joining tree methods are both presented in figure 3 and 4 respectively. It was inferred that the topology of the two phylograms show some level of similarities, as *S. bovis* and *S. curassoni*[18] with *S. intercalatum* being their closest relative, *S. haematobium* and *S. leiperi*, also *S. malayensis* and *S. mekongi* are all closely related as appeared in both phylograms. The result of these phylograms could help our understanding in the pattern of infection of closely related members of the parasite, having considered the phylogenetic relationship in this complex and also attest useful in vector control.




Fig. 1. Nucleic acid composition of the taxa of *Schistosoma* species

* Avg. represent average nucleic acid composition, T (thymine), C (cytosine), A (adenine) and G (Guanine)




Fig. 2. Inferred Ancestral Sequences of the *Schistosoma* species taxa




Fig. 3. Phylogenetic analysis by maximum likelihood and bootstrap percentages (n=3000)




Fig. 4. Phylogenetic analysis by Neighbouring-Joining tree and bootstrap percentages (n=3000)




Fig. 5. An outgroup based phylogenetic analysis by maximum likelihood showing Schistosomes evolutionary trend

Table 1. Sequences of retrieved COI genes in *Schistosoma* species

No.	Species	Accession No.	Size (bp)	Continent
1	<i>S. mansoni</i>	AF130787.1	5,190	Africa
2	<i>S. bovis</i>	AY157212.1	1,125	Africa
3	<i>S. mattheei</i>	AY157211.1	1,125	Africa
4	<i>S. curassoni</i>	AY157210.1	1,125	Africa
5	<i>S. haematobium</i>	AY157209.1	1,125	Africa
6	<i>S. intercalatum</i>	AY157208.1	1,125	Africa
7	<i>S. leiperi</i>	AY157207.1	1,125	Africa
8	<i>S. margrebowiei</i>	AY157206.1	1,125	Africa
9	<i>S. nasale</i>	AY157205.1	1,125	India & S. Asia
10	<i>S. indicum</i>	AY157204.1	1,125	India & S. Asia
11	<i>S. spindale</i>	AY157203.1	1,125	India & S. Asia
12	<i>S. rodhaini</i>	AY157202.1	1,125	Africa
13	<i>S. incognitum</i>	AY157201.1	1,125	Central Asia
14	<i>S. sinensium</i>	AY157197.1	1,125	Central & S. E. Asia
15	<i>S. malayensis</i>	AY157198.1	1,125	Central & S. E. Asia
16	<i>S. mekongi</i>	AY157199.1	1,125	Central & S. E. Asia
17	<i>Schistosoma sp</i>	AJ419784.1	754	-

*S.E- South-East, S- South

Table 2. Maximum Composite Likelihood Estimate of the Pattern of Nucleotide Substitution

	A	T	C	G
A	-	<i>10.94</i>	<i>2.34</i>	<i>16.56</i>
T	<i>5.08</i>	-	<i>3.51</i>	<i>5.02</i>
C	<i>5.08</i>	<i>16.41</i>	-	<i>5.02</i>
G	<i>16.76</i>	<i>10.94</i>	<i>2.34</i>	-

*Each entry shows the probability of substitution (*r*) from one base (row) to another base (column). For simplicity, the sum of *r* values is made equal to 100. Rates of different transitional substitutions are shown in bold and those of transverse substitutions are shown in italics

Table 3. Estimated evolutionary divergence among species

<i>Schistosoma</i> Sp.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>S. mansoni</i>															
<i>S. bovis</i>	0.990														
<i>S. mattheei</i>	1.021	0.135													
<i>S. curassoni</i>	1.012	0.053*	0.137												
<i>S. haematobium</i>	1.001	0.110	0.158	0.113											
<i>S. intercalatum</i>	1.013	0.064*	0.139	0.064*	0.114										
<i>S. leiperi</i>	1.008	0.105	0.137	0.097*	0.103	0.098*									
<i>S. margrebowiei</i>	1.024	0.149	0.170	0.141	0.157	0.148	0.140								
<i>S. nasale</i>	1.019	0.193	0.188	0.194	0.171	0.196	0.181	0.204							
<i>S. indicum</i>	0.969	0.197	0.202	0.197	0.205	0.202	0.189	0.209	0.200						
<i>S. spindale</i>	1.042	0.203	0.206	0.207	0.200	0.210	0.190	0.195	0.194	0.166					
<i>S. rodhaini</i>	0.993	0.218	0.220	0.223	0.203	0.230	0.203	0.237	0.210	0.245	0.219				
<i>S. incognitum</i>	1.065	0.257	0.270	0.251	0.263	0.253	0.258	0.264	0.254	0.253	0.271	0.279			
<i>S. mekongi</i>	1.027	0.260	0.251	0.261	0.241	0.268	0.241	0.265	0.257	0.253	0.266	0.243	0.267		
<i>S. malayensis</i>	1.025	0.256	0.259	0.261	0.254	0.258	0.260	0.262	0.261	0.258	0.256	0.247	0.259	0.067	
<i>S. sinensium</i>	1.102	0.302	0.307	0.312	0.302	0.328	0.311	0.317	0.275	0.292	0.294	0.302	0.313	0.205	0.218

* Areas identified with smallest evolutionary divergence

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fenwick A. Waterborne infetiou diseases- could they be consigned to history? Science. 2006;313:1077-1081.
2. Attwood SW, Fatih FA, Mondal MM, Alim MA, Fadjar S, Rajapakse RP, Rollinson D. A DNA sequence-based study of the *Schistosoma indicum* (Trematoda: Digenea) group: population phylogeny, taxonomy and historical biogeography. Parasitology. 2007;134:2009-2020.
3. Lawton Scott P, Hirohisa Hirai, Joe E Ironside, David A Johnston and David Rollinson. Genomes and geography: genomic insights into the evolution and phylogeography of the genus *Schistosoma*. Parasites & Vectors. 2011;4:131.
4. Agatsuma T. Origin and evolution of *Schistosoma japonicum*. ParasitolInt. 2003;52:335-340.
5. Avise JC. Phylogeography: the history and formation of species. Harvard University press, Cambridge; 2000.
6. Criscione CD, Poulin R, Blouin MS. Molecular ecology of parasites: elucidating ecological and microevolutionary processes. Mol Ecol. 2005;14:2247–2257.
7. Wilkerson RC, Foster PG, Li C, Sallum MA. Molecular phylogeny of neotropical Anopheles (Nyssorhynchus) albitarsis species complex (Diptera: Culicidae), Ann. Entomol. Soc. Am. 2005;98(6):918–925.
8. Khan HA, Arif IA, Bahkali AH, Al Farhan AH, Al Homaidan AA. Bayesian, maximum parsimony and UPGMA models for inferring the phylogenies of antelopes using mitochondrial markers, Evol. Bioinform. 2008;4:263–270.
9. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools, Nucl. Acids Res. 1997;25(24):4876–4882.
10. Benson Otarigho, Mofolusho O. Falade. Molecular Evolution and Phylogenomics of the Anophelesgambiae Complex. Journal of Mosquito Research. 2013;3(9):65-70.
11. Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees, Molecular Biology and Evolution. 1993;10(3):512-526. PMID:8336541
12. Mohanty A, Swain S, Kar SK, Hazra RK. Analysis of the phylogenetic relationship of Anopheles species, subgenus Cellia (Diptera: Culicidae) and using it to define the relationship of morphologically similar species, Infection, Genetics and Evolution. 2009;9(6):1204–1224.
Available: <http://dx.doi.org/10.1016/j.meegid.2009.06.021>. PMID:19577013.
13. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods, Molecular Biology and Evolution. 2011;28(10):2731-2739.

14. Roe AD, Sperling FA. Patterns of evolution of mitochondrial cytochrome c oxidase I and II DNA an implications for DNA barcoding, *Mol. Phylogenet. Evol.* 2007; 44(1):325-345.
Available: <http://dx.doi.org/10.1016/j.ympev.2006.12.005>, PMID:17270468.
15. Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbor-joining method, *Proceedings of the National Academy of Sciences (USA)*. 2004;101(30):11030-11035.
16. Kamali M, Xia A, Tu Z, Sharakhov IV. A New Chromosomal Phylogeny Supports the Repeated Origin of Vectorial Capacity in Malaria Mosquitoes of the *Anopheles gambiae* Complex, *PLoS Pathog.*, 2012;8(10):E1002960, Available: <http://dx.doi.org/10.1371/journal.ppat.1002960>. PMID:23055932 PMCID:3464210.
17. Nei M. and Kumar S. . *Molecular Evolution and Phylogenetics*. Oxford University Press, New York; 2000.
18. Kane RA, Southgate VR, Rollinson D, Littlewood DT, Lockyer AE, Pagès JR, TchuemTchuente LA, Jourdane J. A phylogeny based on three mitochondrial genes supports the division of *Schistosoma intercalatum* into two separate species. *Parasitology*. 2003;127(Pt 2):131-7.
19. Littlewood DT, Lockyer AE, Webster BL, Johnston DA, Le TH: The complete mitochondrial genomes of *Schistosoma haematobium* and *Schistosoma spindale* and the evolutionary history of mitochondrial genome changes among parasitic flatworms. . 2006;39:452-67.
20. Zarowiecki MZ, Huyse T, Littlewood DT: Making the most of mitochondrial genomes- markers for phylogeny, molecular ecology and barcodes in *Schistosoma* (Platyhelminthes: Digenea). *Int J Parasitol* 2007;37:1401-1148.
21. *Schistosoma haematobium*. IARC Monograph- 100B: 2012; pg 371-384.

© 2014 Onile et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sciencedomain.org/review-history.php?iid=419&id=32&aid=3583>