



# Knowledge and Practice of Antimicrobial use for Treatment of Urinary Tract Infections among Female Medical Students in Enugu, Nigeria: A Cross-sectional Study

Chikaodili Erica Odoh <sup>a\*</sup>, Iheoma Amara Ochiobi <sup>a</sup>,  
Chukwuebuka Obiakalusi <sup>a</sup>, Linda Uzo Amara <sup>a</sup>  
and Chika N. Onwasigwe <sup>a,b,c</sup>

<sup>a</sup> Department of Community Medicine, Enugu State University (ESUT) Teaching Hospital, Parklane, Enugu, Nigeria.

<sup>b</sup> Department of Community Medicine, College of Medicine, University of Nigeria Enugu Campus, Enugu, Nigeria.

<sup>c</sup> Institute of Public Health, University of Nigeria Nsukka (Enugu Campus), Enugu, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors CEO, IAO, CO, LUA and CNO contributed to the conceptualization and design of the study, methodology, data acquisition, formal analysis, interpretation of data, wrote, reviewed and edited the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijtdh/2024/v45i121609>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127589>

**Short Research Article**

**Received: 27/09/2024**

**Accepted: 29/11/2024**

**Published: 04/12/2024**

\*Corresponding author: Email: [ericachikaodiliodoh@gmail.com](mailto:ericachikaodiliodoh@gmail.com);

## ABSTRACT

**Background:** Urinary tract infections (UTI) are a significant clinical problem empirically treated with antibiotics that could worsen the spread of antibiotics resistance.

**Objective:** We described the knowledge, attitude, and treatment practices of UTI among female medical students in Enugu, southeast Nigeria.

**Methods:** A cross-sectional, questionnaire-based survey involving female medical students at Enugu State University of Technology, in Enugu, Nigeria was conducted in October 2023. Overall, 180 students were enrolled. The questionnaires were constructed after systematically reviewing the literature. The data were analyzed by calculating frequencies and means.

**Results:** Our study sample includes 174 female medical students (Response rate, 96.7%). Most of the students are aged 20–24 years (63.2%), with mean (standard deviation) of 23.5 ( $\pm$  4.6) years, single (86.8%), Igbo (97.1%), and Christian (97.7%). Most students (166 students, 95.4%) had adequate knowledge of UTI, but a minority of students (45.5%) sought treatment from a trained healthcare provider for proper laboratory investigation, diagnosis, and treatment of their previous UTI episodes. Among those that did not seek care with trained healthcare providers, most, 27 (75.0%), self-treated. Most students (27 students, 90.0%) positively practiced completion of their prescribed drugs. The most commonly drugs used for self-medication of UTI were Amoxicillin, Canesten cream, Ciprofloxacin, and Gentamycin.

**Conclusion:** Although our study shows female medical students have adequate knowledge about UTIs, their treatment practices reveal that strengthening educational efforts on antibiotic stewardship is essential to ensure that medical students not only understand the theoretical aspects of UTIs but also apply best practices in their treatment. Our study suggests the need for improved education on antibiotics stewardship to address the spread of antimicrobial resistance, and ensuring effective treatment of UTIs in the future.

**Keywords:** Antibiotics resistance; knowledge; attitudes; practice; urinary tract infections; medical students; Nigeria.

## 1. INTRODUCTION

Urinary tract infections (UTIs) occur when microbial pathogens invade the urinary tract, causing symptoms. Cystitis affects the lower urinary tract, while pyelonephritis affects the upper tract (Gupta et al., 2017; John Mwang and Innocent Mchami, 2022; Foxma, 2014; Flores-Mireles et al., 2015 ). UTIs are one of the common bacterial infections seen in primary health care, second only to infections of the respiratory tract (Foxman, 2014). UTIs are common bacterial infections, especially among women, with an incidence of about 1% in school-aged girls and 4% in women of childbearing age (John Mwang and Innocent Mchami, 2022; Foxman, 2014). The incidence increases with sexual activity and childbearing, reaching 25-30% in women aged 20-40 (John Mwang and Innocent Mchami, 2022). *Escherichia coli* is the most common cause of uncomplicated UTIs, but other pathogens include *Enterococcus faecalis*, *Enterobacter species*, *Staphylococcus saprophyticus*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas species*, fungi (candidiasis), and protozoa (*Trichomonas vaginalis*) (John Mwang and Innocent Mchami,

2022; Flores-Mireles et al., 2015; Alshahrani et al., 2022).

UTIs are caused by pathogenic invasion of the urothelium, leading to inflammation in both the upper and lower urinary tracts (Foxman, 2014). UTIs are the most common bacterial infections in humans and are frequently seen in hospitals, accounting for about 40% of all nosocomial bacterial infections worldwide (Foxman, 2014). They are a leading cause of morbidity and mortality, particularly in the first two years of life, in women, and in the elderly, with significant financial implications (Kolawole et al., 2009). UTIs are also a common reason for outpatient visits, making up over 20% of such visits to general practitioners (Gupta et al., 2017). Epidemiological studies show that about 33% of women will experience at least one UTI in their lifetime, requiring antibiotic treatment (Foxman, 2014). Recurrence within six months of the initial infection occurs in 20-50% of cases (Flores-Mireles et al., 2015). UTIs significantly impact quality of life, affecting social, relational, and occupational activities.

UTIs can be classified based on severity levels, presence of risk factors, and microbiological

classification of pathogens. Clinically, UTIs can present as cystitis, pyelonephritis, or urosepsis (Flores-Mireles et al., 2015). Asymptomatic bacteriuria, while not an infection, is a risk factor for UTIs in certain situations. The severity of UTIs varies: Cystitis is less severe than pyelonephritis, which can range from mild to severe. Urosepsis is the most severe form, with severity graded similarly to sepsis: sepsis, severe sepsis, and septic shock. UTIs are also categorized as: uncomplicated which occurs in healthy individuals with normal urinary tracts; and complicated involve abnormal urinary tract anatomy or function, concomitant diseases, or failed previous treatments.

Symptoms in adults include dysuria, urinary urgency, frequency, nocturia, painful voiding, bladder discomfort, and stranguria (John Mwang and Innocent Mchami, 2022; Foxman, 2014; Flores-Mireles et al., 2015). Pain at the start or during urination suggests a urethral issue, while post-voiding pain indicates bladder or prostate problems (Gupta et al., 2017; Foxman, 2014). Women are more prone to UTIs due to their shorter urethra and proximity to the anus, along with factors like inadequate bladder emptying, instrumentation, urinary tract calculi, indwelling catheters, urinary obstruction, contraceptive use, anal intercourse, constipation, pregnancy, and immunosuppression (Gupta et al. 2017, Flores-Mireles et al, 2015; Esan et al., 2023; Dessie et al., 2024).

Women are at higher risk for UTIs due to their shorter urethra and certain behaviours, such as delayed micturition, sexual activity, and use of diaphragms and spermicides (Flores-Mireles et al., 2015). Infections often result from bacteria entering the urethra and ascending to the bladder, commonly linked to sexual activity or catheterization (Foxman, 2014). Postmenopausal women are also at higher risk due to factors like bladder or uterine prolapse, loss of oestrogen, and higher likelihood of medical conditions like diabetes (Flores-Mireles et al., 2015). Studies indicate a seasonal peak in UTI incidence, particularly in the post-summer months. Antibiotics are essential for treating symptomatic UTIs, with choices based on pathogen susceptibility, efficacy, resistance potential, safety, side effects, cost, and availability (Flores-Mireles et al., 2015). Recommended regimens for uncomplicated cystitis and pyelonephritis in premenopausal women include: Fosfomycin trometamol, Nitrofurantoin, Pivmecillinam, and Trimethoprim/sulfamethoxazole. Empiric

treatment for uncomplicated UTIs often starts with trimethoprim/sulfamethoxazole in communities with low resistance rates. Penicillin use is significantly associated with multi-drug-resistant Enterobacteriaceae UTI (Rossignol et al. 2015). In resistant areas, fluoroquinolones like ciprofloxacin, levofloxacin, norfloxacin, and ofloxacin are used (Flores-Mireles et al., 2015).

Diagnosing a UTI requires laboratory tests on urine samples. Urine microscopy can reveal red and white blood cells, indicating infection (John Mwang and Innocent Mchami, 2022; Flores-Mireles et al., 2015). A urine culture is positive if it shows a bacterial colony count of at least 10,000 colony-forming units per ml. These cultures also help determine antibiotic sensitivity, guiding treatment choices (Flores-Mireles et al., 2015). Women with recurrent UTIs can often self-diagnose based on symptoms, with an 84% accuracy rate. Positive predictors for recurrent UTIs include symptoms after intercourse, a history of pyelonephritis, absence of nocturia, and quick symptom resolution after treatment. Negative predictors include nocturia and persistent symptoms between infections. Urinalysis is essential for patients with dysuria, with the gold standard being a spun midstream clean-catch urine specimen. Bacteria or pyuria (or both) are typically present in UTI cases. Leukocyte esterase is 75% sensitive and 98% specific for UTI detection, while positive nitrite is 90% specific but only 30% sensitive. Urine culture is generally unnecessary for patients with consistent symptoms and a positive dipstick test unless there are risk factors for complicated infections (Gupta et al., 2017; Flores-Mireles et al., 2015).

Inappropriate antimicrobial use is common among female university students, with studies showing a tendency towards self-medication, especially among pharmacy students (Ibrahim Sharif and Suleiman, 2013). This behaviour can significantly impact future antimicrobial use, particularly for UTI treatment. The misuse of antibiotics is a global issue, prevalent in both developing and developed countries, including Pakistan, Nepal, Eritrea, Uzbekistan, Oman, Sudan, Jordan, Lebanon, Yemen, and Nigeria (Naber et al., 2009). Commonly used antibiotics for UTIs include Ampicillin, Amoxicillin, Cefotaxime, Ceftriaxone, Ciprofloxacin, Gentamicin, Cefoxitin, Erythromycin, Tetracycline, and Vancomycin. Recommended regimens for acute uncomplicated cystitis and pyelonephritis include short-course treatments

(single dose to 5 days), which are as effective as longer regimens with fewer side effects (Flores et al., 2015; Naber et al., 2009).

Antimicrobials are crucial for treating infections, including UTIs (Gupta et al., 2017; Flores et al., 2015). However, antimicrobial resistance (AMR) is a significant challenge, especially in developing countries where antibiotics are easily accessible without prescriptions (Flores et al., 2015). Inappropriate use of antimicrobials, such as self-medication, can lead to harmful effects like diarrhoea and skin rashes and exacerbate AMR (Gupta et al., 2017; Flores et al., 2015). Global actions to combat AMR include improving awareness, understanding misuse effects, communication, education, and training (Leung et al., 2011). This global action plan to tackle AMR is strongly linked with the drive towards universal health coverage (UHC). Tackling AMR is crucial for maintaining the effectiveness of antibiotics, which are essential for treating infections and enabling other medical procedures. This directly supports the goals of UHC by ensuring that effective treatments remain available and affordable for all (Eze et al., 2024; Bloom et al., 2017; Joshua et al., 2021).

This study focuses on the knowledge, attitude, and practice of inappropriate antibiotic use for UTIs among female medical students, highlighting the need for better awareness and management. Our study findings could inform health policy on interventions to tackle AMR in the region.

## 2. METHODOLOGY

### 2.1 Study Design

This descriptive cross-sectional study was conducted in October 2023, focusing on female medical students at Enugu State University Teaching Hospital. All eligible female medical students were included, following the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines to ensure proper reporting of the study's design, conduct, and findings (Von Elm et al., 2007).

### 2.2 Study Setting

The study took place at ESUT Teaching Hospital, located in Parklane GRA, Enugu, the capital of Enugu State, Nigeria. Enugu State, with an estimated population of 5.2 million in 2023, is known for its coal deposits and is often referred to as the 'Coal City State'. It features highlands in

Awgu, Udi, and Nsukka to the east and the Oji River basin to the west. The economy is primarily rural and agrarian, with significant trading and civil service sectors (Lawani et al., 2022; Eze et al., 2022). The population is predominantly Christian, with a minority practicing Islam and African traditional religions. The state has two main tertiary health centres: ESUT Teaching Hospital and the University of Nigeria Teaching Hospital (UNTH), along with several private and missionary primary and secondary healthcare centres (Lawani et al.; 2022).

ESUT Teaching Hospital has approximately 1,061 students, though the exact number fluctuates due to admissions, transfers, and suspensions. ESUT Teaching Hospital has three major hostels for medical students, while those studying Anatomy and Medical Laboratory Science live off-campus. The student body is mostly Igbo, and the primary languages spoken are Igbo and English.

### 2.3 Study Participants

All female medical students at ESUT Teaching Hospital, regardless of age, class, or residence, were eligible to participate. Women are more susceptible to UTIs because of their shorter urethra, prolonged delaying urination, engaging in sexual activity, and using diaphragms and spermicides (Gupta et al., 2017; Flores et al., 2015). Exclusions included female anatomy and medical laboratory science students and those who opted out of the study.

### 2.4 Sample Size

Based on an 82% prevalence rate from a previous study (Mangai et al., 2019), the sample size was calculated using a design effect (DEFF) of 1.05, a 5% significance level, a 5% margin of error, and a 10% inflation to account for non-response. The minimum estimated sample size was 180.

### 2.5 Sampling Procedure

A simple random sampling technique was used to select participants. Stratified sampling ensured representation across all departments and levels within the university. To avoid bias, participants were not informed in advance. The purpose of the questionnaire was explained, and participation was voluntary. Questionnaires were distributed randomly to students in class, collected after 30 minutes, and followed by an educational session on UTIs, their diagnosis, and antibiotic treatment.

## 2.6 Study Instrument

Data were collected using a self-administered structured questionnaire, distributed to students in their respective departments. The questionnaire was divided into four sections. Section A: Socio-demographic variables (age, marital status, religion, ethnicity, year of study). Section B: Knowledge variables (awareness and knowledge of UTI, initial source of information, definition of UTI). Section C: Attitudinal variables (history of UTI, symptoms, diagnosis). Section D: Practice variables (hospital visits for diagnosis and treatment, completion of prescribed dosage, alternative treatments, and drugs used).

## 2.7 Data Analysis and Presentation

Simple frequency tables with percentages were produced using IBM SPSS®.

## 3. RESULTS

### 3.1 Sociodemographic Characteristics of Study Participants

The most common age group among the study participants was 20-24 years (63.2%), with a mean age of 23.5 years ( $\pm 4.6$ ) – Table 1. Most participants were single (86.8%), Igbo (97.1%), and Catholic (58.6%). A significant proportion were in their fifth year of study (44.3%) – Table 1.

### 3.2 Knowledge of UTI Among Study Participants

Most participants had knowledge of UTIs (97.1%), with the majority learning about the symptoms and treatment from school (64.4%) – Table 2. The majority accurately identified that a UTI is an infection of the urethra, bladder, ureter, and kidney (95.4%), and most had adequate knowledge of UTI symptoms.

**Table 1. Sociodemographic characteristics of study participants**

	Frequency	Percent
<b>Age</b>		
◦ < 20 years	16	9.2%
◦ 20 – 24 years	110	63.2%
◦ 25 – 29 years	34	19.5%
◦ 30 – 34 years	9	5.2%
◦ 35 – 39 years	2	1.1%
◦ ≥ 40 years	3	1.7%
<b>Marital status</b>		
◦ Single	151	86.8%
◦ Married	22	12.6%
◦ Divorced/Widowed	1	0.6%
<b>Religion</b>		
◦ Catholic	102	58.6%
◦ Pentecostal/Protestants	68	39.1%
◦ Muslim	1	0.6%
◦ Africa Traditional Religion	3	1.7%
<b>Ethnicity</b>		
◦ Igbo	169	97.1%
◦ Yoruba	4	2.3%
◦ Others	1	0.6%
<b>Year of study</b>		
◦ Second year	14	8.0%
◦ Third year	18	10.3%
◦ Fourth year	38	21.8%
◦ Fifth year	77	44.3%
◦ Sixth year	27	15.5%

**Table 2. Knowledge of UTI among female medical students**

	Frequency	Percent
<b>Ever heard of UTI</b>		
◦ No	5	2.9%
◦ Yes	169	97.1%
<b>First source of information</b>		
◦ Radio	4	2.3%
◦ Television	9	5.2%
◦ Newspaper	2	1.1%
◦ School	112	64.4%
◦ Relatives	15	8.6%
◦ Friends	13	7.5%
◦ Spouse	1	0.6%
◦ Church	1	0.6%
◦ Hospital/Clinic	9	5.2%
◦ Others	8	4.6%
<b>Knowledge of what UTI is</b>		
◦ UTI is an airborne disease: Correct answer = No	172	98.9%
◦ UTI is an infection of the liver: Correct answer = No	171	98.3%
◦ UTI is an infection of the heart: Correct answer = No	168	96.6%
◦ UTI is an infection urethra, bladder, ureter, and/or kidney: Correct answer = Yes	166	95.4%
◦ UTI is transmitted through sexual intercourse: Correct answer = No	167	96.0%
<b>Specific knowledge of UTI symptoms</b>		
◦ Pain urination	137	78.7%
◦ Fever	99	56.9%
◦ Inability to hold urine	87	50.0%
◦ Discharge from urinary tract	125	71.8%
◦ Frequent urination	102	58.6%
◦ Having urgent urge to urinate	96	55.2%

### 3.3 Attitude of Study Participants Towards UTI

Fig. 1 shows a Likert chart illustrating the attitudes of study participants towards UTIs. Generally, participants had a positive attitude towards using antimicrobials for UTI treatment.

### 3.4 Practice of Study Participants Towards UTI

Most participants had never experienced a UTI (62.1%) – Table 3. Among those who had, most did not seek medical treatment from a trained healthcare provider (54.5%). Of those who received a prescription from a trained healthcare provider, most completed their prescribed medication (90.0%). Among participants with a history of UTIs, most self-treated with a cocktail

of antibiotics: Septrin, Ampiclox, Amoxicillin, Canesten cream, Ciproxin, and Gentamycin (75.0%) – Table 3. Those who did not seek treatment from a trained healthcare provider often cited financial constraints as the reason – Table 4.

## 4. DISCUSSION

Our study shows that the majority of female medical studies have good knowledge of UTI, with most of them receiving their first information from school. Our study contrasts with a study of healthcare students at the University of Science and Technology, Ajman, UAE, which found 92.3% of students using antimicrobials irrationally (Ibrahim et al., 2013). In our study, 54.5% of students used antimicrobials inappropriately, indicating a significant decline in




Fig. 1. Attitude of female medical students in Enugu State University Teaching Hospital towards UTI, 2023

**Table 3. Practice of UTI among female medical students**

	Frequency	Percent
<b>Ever suffered from UTI</b>		
◦ No	108	62.1%
◦ Yes	66	37.9%
<b>Sought treatment in a hospital</b>		
	n = 66	
◦ No	36	54.5%
◦ Yes	30	45.5%
<b>Completed prescribed medications</b>		
	n = 30	
◦ No	3	10.0%
◦ Yes	27	90.0%
<b>What treatment did you get?</b>		
	n = 36	
◦ I treated myself	27	75.0%
◦ Traditional healer	0	0.0%
◦ Patent medicine vendor (Chemist)	7	19.4%
◦ Ignored the symptoms	2	5.6%
<b>Drugs used for self-medication</b>		
	n = 27	
◦ Septrin	1	3.7%
◦ Ampiclox	3	11.1%
◦ Amoxicillin	9	33.3%
◦ Canesten cream	7	25.9%
◦ Ciproxin	17	63.0%
◦ Gentamycin injection	4	14.8%
◦ Others	1	3.7%

**Table 4. Reasons for not seeking care with trained healthcare providers, n = 36**

Reasons for not seeking care	No	Yes
◦ Against my faith	35 (97.2%)	1 (2.8%)
◦ I do not have money	34 (94.4%)	2 (5.6%)
◦ I do not have time	23 (63.9%)	13 (36.1%)
◦ I am shy to talk about it	32 (88.9%)	4 (11.1%)
◦ I believe it was sexually transmitted	33 (91.7%)	3 (8.3%)

misuse. In a survey at a urinary tract infection/AIDS clinic at Konfo Anokye Teaching Hospital in Kumasi, Ghana, 74.5% of patients admitted to self-medicating before visiting the clinic, similar to the 75.0% in our study (Boaitey et al. 2012). However, the prevalence of self-medication in our study is higher than the prevalence reported in a similar study in Bahrain which reported a prevalence of 44.8% (James et al., 2006).

This difference may be attributed to our academic environment, where there is greater awareness of drug use and its harmful side effects. The irrational use of antimicrobials among female undergraduates is common, often due to economic factors, time constraints, and the desire to take responsibility for one's health (Alshahrani et al., 2022; Mangai et al., 2019). However, this practice can lead to hazardous

consequences such as misdiagnosis (Gupta et al., 2017; Flores et al., 2015). The problem is more prevalent in developing countries where antimicrobials are available without prescription (John and Innocent, 2022; Joshua et al., 2021; Torres et al., 2019). The prevalence of self-medication in our study is comparable to that reported among female students in Egypt, Ethiopia, India, Saudi Arabia, and Uganda, but higher than in many European countries and lower than among Palestinian female students (Alshahrani et al., 2022; Dessie et al., 2024; Abdelbadee et al., 2024; Nalwadda and Were, 2024; Jelly et al., 2022).

Factors contributing to the high prevalence of self-medication include lack of time to visit a clinic, high economic status, overconfidence among students, prescribing patterns of general practitioners, and liberal dispensing by

pharmacists (Torres et al., 2019; Araia et al., 2019; Ahmadi et al., 2016; Abdi et al., 2018; Ezeet al., 2022; Lukovic et al., 2014). This is alarming given the high awareness of microbial resistance associated with misuse (Leung et al., 2011; Bloom et al., 2017; Joshua et al., 2021). The prevalence of self-medication in our study (75.0%) is comparable to the 79.2% prevalence reported in Eritrea (Araia et al., 2019), 79.9% reported in Belgrade, Serbia (Lukovic et al., 2014), and 89.6% prevalence reported in Kermanshah, Iran (Abdi et al., 2018). The prevalence of self-medication in our study, however, is significantly higher than the prevalence of 33.7% reported in another study in Kermanshah, Iran (Ahmadi et al., 2016). More than one-third of Palestinian students admitted not completing their prescribed dosage, higher than 10.0% of ESUT Teaching Hospital students (Sawalha, 2008). Our findings suggest that liberal dispensing by pharmacists contributes to antimicrobial misuse (Torres et al., 2019). Incorporating topics on the rational use of drugs and prescription writing into health science curricula could reduce inappropriate use (Joshua et al., 2021; Mangaiet al., 2019; Eze et al., 2020). Efforts by health authorities and academic circles to increase public awareness are essential (Joshua et al., 2021).

The most commonly self-used antimicrobials were Ciproxin, Amoxicillin, Canesten cream, and Ampiclox, likely due to frequent prescribing by physicians and liberal dispensing by pharmacists (Torres et al., 2019). Some students (19.4%) relied on advice from pharmacists and patent medicine dealers, highlighting the important roles of these professionals in promoting responsible self-medication (Araia et al., 2019).

#### **4.1 Study Limitations**

Our study has a few limitations. First, we acknowledge that the findings rely on self-reported data, which may be influenced by recall and response biases. Secondly, the cross-sectional nature of the study restricts our capacity to determine causality. Finally, we did not employ recent empirical approach such as regression or matching in our study limiting our ability to draw firm conclusions.

#### **4.2 Implications for Public Health Practice and Research**

Based on our findings, we recommend a multifaceted approach to address the misuse of

antimicrobials and improve the diagnosis and treatment of UTIs. This includes disseminating information on the hazards of antimicrobial misuse and the importance of proper diagnosis and treatment through various channels such as health talks, seminars, flyers, student forums, and media programs. Additionally, we advocate for incorporating health education on the irrational use of antimicrobials at all educational levels in Nigeria to foster a better understanding from an early age. Furthermore, we suggest monitoring the activities of medicine dealers to ensure they provide accurate information and guidance, thereby increasing public knowledge on the proper approach to diagnosing and managing UTIs. This comprehensive strategy aims to enhance awareness, education, and regulation to combat antimicrobial resistance effectively (Joshua et al., 2021, Parveen et al., 2022; Roca et al., 2015; Almaghlouth et al., 2023).

### **5. CONCLUSION**

In conclusion, while many students are aware of UTIs, misconceptions persist, and some students resort to self-medication or patent medicine dealers instead of seeking medical attention. This leads to undiagnosed cases and partial treatments. There is a lack of knowledge about the long-term complications of UTIs and the adverse effects of self-medication. Improving educational efforts on antibiotic stewardship is essential to ensure that medical students not only understand the theoretical aspects of UTIs but also apply best practices in their treatment. This approach is vital for curbing the spread of antibiotic resistance and ensuring effective treatment of UTIs in the future.

### **DATA AVAILABILITY STATEMENT**

The datasets analyzed in this study are available from the corresponding author on reasonable request.

### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

### **ETHICS APPROVAL AND CONSENT**

Permission was obtained from ESUT Teaching Hospital. The aim and nature of the study was

also explained to the participants and Informed consent obtained prior to administration of the questionnaire.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Abdallah, S. R., Abd-Elazim, M. H., Hasab Allah, M. F., & Soliman, H. A. (2024). Knowledge and habitual practice regarding prevention of genito-urinary tract infection among adolescent nursing girls students. *Minia Scientific Nursing Journal*, 15(2), 21–30.
- Abdi, A., Faraji, A., Dehghan, F., & Khatony, A. (2018). Prevalence of self-medication practice among health sciences students in Kermanshah, Iran. *BMC Pharmacology and Toxicology*, 19, 36.
- Ahmadi, S. M., Jamshidi, K., Sadeghi, K., Abdi, A., & Vahid, M. P. (2016). The prevalence and affecting factors on self-medication among students of Kermanshah University of Medical Science in 2014. *Journal of Clinical and Diagnostic Research*, 10(5), IC01–IC04.
- Almaghlouth, A. K., Alkhalaf, R. A., Alshamrani, A. A., Albrahim, J. A., Alhulibi, B. S., Al-Yousef, A. Y., et al. (2023). Awareness, knowledge, and attitude towards urinary tract infections: An appraisal from Saudi Arabia. *Cureus*, 15(11), e49352. DOI: 10.7759/cureus.49352
- Alshahrani, M., Bakheet, S. A., Alzahrani, A., Alzahrani, A., Alqhtani, A. A., Alwabel, H., & Asiri, K. M. (2022). Knowledge, attitude, and practice of urinary tract infection among females in Aseer region. *Bahrain Medical Bulletin*, 44(1), 769–774.
- Araia, Z. Z., Gebregziabher, N. K., & Mesfun, A. B. (2019). Self-medication practice and associated factors among students of Asmara College of Health Sciences, Eritrea: A cross-sectional study. *Journal of Pharmaceutical Policy and Practice*, 12, 3. Available from: <https://doi.org/10.1186/s40545-019-0165-2>
- Bloom, G., Merrett, G. B., Wilkinson, A., Lin, V., & Paulin, S. (2017). Antimicrobial resistance and universal health coverage. *BMJ Global Health*, 2(4), e000518.
- Boaitey, Y. A., Nkrumah, B., Idriss, A., & Tay, S. C. K. (2012). Gastrointestinal and urinary tract pathogenic infections among HIV seropositive patients at the Komfo Anokye Teaching Hospital in Ghana. *BMC Research Notes*, 5, 454.
- Dessie, Z., Gela, D., & Yusuf, N. T. (2024). Knowledge, attitude, practice, and associated factors toward the prevention of catheter-associated urinary tract infection among nurses in Ethiopia: Cross-sectional study. *SAGE Open Nursing*, 10, 1–10. Available from: <https://doi.org/10.1177/23779608241281317>
- Esan, D. T., Sokan-Adeaga, A. A., Bello, C. B., Olabisi, O. I., Odugbume, B., & Ajayi, P. O. (2023). Knowledge, attitude, and practice of antenatal mothers toward urinary tract infection in selected health facilities in Ekiti State, Nigeria. *Journal of Public Health Research*, 12(3), 1–12. Available from: <https://doi.org/10.1177/22799036231197180>
- Eze, P., Aniebo, C. L., Agu, U. J., Agu, S. A., & Acharya, Y. (2022). Validity of maternal recall for estimating childhood vaccination coverage – Evidence from Nigeria. *Vaccine*, 40(1), 28–36. Retrieved from <https://doi.org/10.1016/j.vaccine.2021.11.060>
- Eze, P., Idemili, C. J., & Lawani, L. O. (2024). Evaluating health systems' efficiency towards universal health coverage: A data envelopment analysis. *Inquiry*, 61, 1–12. Retrieved from <https://doi.org/10.1177/00469580241235759>
- Eze, P., Lawani, L. O., Agu, U. J., & Acharya, Y. (2022). Catastrophic health expenditure in sub-Saharan Africa: Systematic review and meta-analysis. *Bulletin of the World Health Organization*, 100(5), 337–351. Available from: <http://dx.doi.org/10.2471/BLT.21.287673>
- Eze, P., Lawani, L. O., Chikezie, R. U., Ukaegbe, C. I., & Iyoke, C. A. (2020). Perinatal outcomes of babies delivered by second-stage Caesarean section versus vacuum extraction in a resource-poor setting, Nigeria – A retrospective analysis. *BMC Pregnancy and Childbirth*, 20, 298. Available from: <https://doi.org/10.1186/s12884-020-02995-9>
- Flores-Mireles, A. L., Walker, J. N., Caparon, M., & Hultgren, S. J. (2015). Urinary tract infections: Epidemiology, mechanisms of infection and treatment options. *Nature Reviews Microbiology*, 13(5), 269–284.

- Foxman, B. (2014). Urinary tract infection syndromes: Occurrence, recurrence, bacteriology, risk factors, and disease burden. *Infectious Diseases Clinics of North America*, 28(1), 1–13.
- Gupta, K., Grigoryan, L., & Trautner, B. (2017). Urinary tract infection. *Annals of Internal Medicine*, 167(7), ITC49–64.
- Ibrahim Sharif, S., & Suleiman Sharif, R. (2013). Antibiotics use with and without a prescription in healthcare students. *American Journal of Pharmacological Sciences*, 1(5), 96–99.
- James, H., Handu, S. S., Al Khaja, K. A. J., Otoom, S., & Sequeira, R. P. (2006). Evaluation of the knowledge, attitude and practice of self-medication among first-year medical students. *Medical Principles and Practice*, 15(4), 270–275.
- Jelly, P., Verma, R., Kumawat, R., Choudhary, S., Chadha, L., & Sharma, R. (2022). Occurrence of urinary tract infection and preventive strategies practiced by female students at a tertiary care teaching institution. *Journal of Education and Health Promotion*, 11(1), 122.
- Joshua, I. A., Bauche, J., & Abdulla, S. (2021). Managing antimicrobial resistance from medical and veterinary health systems perspectives to achieving universal health coverage in the African region. *South East European Journal of Public Health*, 2021(Special Issue 2).
- Kolawole, A. S., Kolawole, O. M., Kandaki-Olukemi, Y. T., Babatunde, S., Durowade, K. S., & Kolawole, C. F. (2009). Prevalence of urinary tract infections (UTI) among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State, Nigeria. *International Journal of Medicine and Medical Sciences*, 1(5), 163–167. Available from: <https://www.researchgate.net/publication/236649983>
- Lawani, L. O., Enebe, J. T., Eze, P., Igboke, F. N., Ukaegbe, C. I., Ugwu, M. O., et al. (2022). Interpregnancy interval after a miscarriage and obstetric outcomes in the subsequent pregnancy in a low-income setting, Nigeria: A cohort study. *SAGE Open Medicine*, 10, 1–11.
- Leung, E., Weil, D. E., Raviglione, M., & Nakatani, H. (2011). The WHO policy package to combat antimicrobial resistance. *Bulletin of the World Health Organization*, 89(5), 390–392.
- Lukovic, J. A., Miletic, V., Pekmezovic, T., Trajkovic, G., Ratkovic, N., Aleksic, D., et al. (2014). Self-medication practices and risk factors for self-medication among medical students in Belgrade, Serbia. *PLOS One*, 9(12), e114644.
- Mangai, M. J., Gaknung, B., Hosea, G., Peter, U., Patience, K., Shikpup, N., et al. (2019). Assessment of knowledge and prevention practices of urinary tract infection (UTI) among female students residing in the University of Jos. *International Research Journal of Public and Environmental Health*, 6(5), 89–96.
- Mwang, J., & Mchami, J. I. (2022). The aetiology and prevalence of urinary tract infections in Sub-Saharan Africa: A systematic review. *Journal of Health & Biological Sciences*, 10(1), 1–7.
- Naber, K. G., Cho, Y. H., Matsumoto, T., & Schaeffer, A. J. (2009). Immunoactive prophylaxis of recurrent urinary tract infections: A meta-analysis. *International Journal of Antimicrobial Agents*, 33(2), 111–119.
- Nalwadda, R., & Were, A. (2024). Knowledge, attitude, and practice towards the prevention of urinary tract infections among female students at Kampala School of Health Sciences, Wakiso: A cross-sectional study. *SJ Obstetrics and Gynecology Africa*, 1(8), 1–10. Available from: <https://doi.org/10.51168/ydp7mt40>
- Parveen, S., Garzon-Orjuela, N., Amin, D., McHugh, P., & Vellinga, A. (2022). Public health interventions to improve antimicrobial resistance awareness and behavioural change associated with antimicrobial use: A systematic review exploring the use of social media. *Antibiotics*, 11(5), 669. Available from: <https://doi.org/10.3390/antibiotics11050669>
- Roca, I., Akova, M., Baquero, F., Carlet, J., Cavalieri, M., Coenen, S., et al. (2015). The global threat of antimicrobial resistance: Science for intervention. *New Microbes and New Infections*, 6, 22–29.
- Rosignol, L., Maugat, S., Blake, A., Vaux, S., Heym, B., Le Strat, Y., et al. (2015). Risk factors for resistance in urinary tract infections in women in general practice: A cross-sectional survey. *Journal of Infection*, 71(3), 302–311. Available from: <https://doi.org/10.1016/j.jinf.2015.05.012>
- Sawalha, A. F. (2008). A descriptive study of self-medication practices among Palestinian medical and nonmedical university

- students. *Research in Social and Administrative Pharmacy*, 4(2), 164–172.
- Torres, N. F., Chibi, B., Middleton, L. E., Solomon, V. P., & Mashamba-Thompson, T. P. (2019). Evidence of factors influencing self-medication with antibiotics in low and middle-income countries: A systematic scoping review. *Public Health*, 168, 92–101.
- Von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., & Vandenbroucke, J. P. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting observational studies. *Bulletin of the World Health Organization*, 85(11), 867–872.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/127589>